

Rabbit IgG Isotype Control

<u>Cat. No.</u>	<u>Form</u>	<u>Quantity</u>
0111-01	Purified (UNLB) IgG	10 mg
0111-02	Fluorescein (FITC) Conjugate	100 tests
0111-09	R-phycoerythrin (PE) Conjugate	100 tests
0111-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg

DESCRIPTION

Source Normal rabbit serum
Ig Isotype Rabbit IgG
Specificity N/A

RESEARCH APPLICATIONS

- Flow cytometry
- Enzyme-Linked-Immunosorbent-Assay (ELISA)
- Fluorescent-Linked-Immunosorbent-Assay (FLISA)

CHARACTERIZATION

To insure lot-to-lot consistency, each batch of product is tested by ELISA, FLISA, and/or flow cytometry for conformance to characteristics of a standard reference reagent.

WORKING DILUTIONS

Flow Cytometry:

Purified IgG	$\leq 1 \mu\text{g}/10^6$ cells
Fluorescein conjugate	$10 \mu\text{L}/10^6$ cells
R-phycoerythrin conjugate	$10 \mu\text{L}/10^6$ cells

Other Applications: Since applications vary, each investigator should determine the optimum working dilutions of the product that is appropriate for their specific needs.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

HANDLING AND STORAGE

- The purified (UNLB) IgG is supplied as 10 mg of purified immunoglobulin in 2.0 mL of 100 mM borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 100 tests in 1.0 mL PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 100 tests in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- With the exception of the low endotoxin, azide-free product, the vials should be stored at 2-8°C. Conjugated forms should **not** be frozen and should be protected from prolonged exposure to light. Aliquot and freeze the low endotoxin, azide-free product at -20°C immediately upon receipt. Each reagent is stable for the period shown on the bottle label if stored as directed.

WARNING

Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.