

## Goat Anti-Hamster IgG (H+L) Absorbed against mouse & rat immunoglobulins

Cat. No.	Form	Quantity
6061-01	Purified (UNLB) Antibody	1.0 mg
6061-02	Fluorescein (FITC) Conjugate	1.0 mg
6061-04	Alkaline Phosphatase (AP) Conjugate	1.0 mL
6061-05	Horseradish Peroxidase (HRP) Conjugate	1.0 mL
6061-08	Biotin (BIOT) Conjugate	1.0 mg
6061-09	R-phycoerythrin (R-PE) Conjugate	0.5 mg
6061-11	Allophycocyanin (APC) Conjugate	0.5 mg
6061-17	*R-phycoerythrin-Cyanine 7 (R-PE-CY™7) Conjugate	0.25 mg

### DESCRIPTION

**Source:** Pooled antisera from goats hyperimmunized with Golden Syrian hamster IgG.  
**Cross Absorption:** Mouse and rat immunoglobulins  
**Purification:** Affinity chromatography on hamster IgG covalently linked to agarose.  
**Specificity:** Reacts with the heavy and light chains of Syrian hamster IgG as demonstrated by ELISA, FLISA, and/or flow cytometry. May also react with the light chains of other hamster immunoglobulins. Minimal cross reactivity with mouse and rat immunoglobulins

### RESEARCH APPLICATIONS

- Immunofluorescent staining
- Enzyme-Linked-Immunosorbent-Assay (ELISA)
- Fluorescent-Linked-Immunosorbent-Assay (FLISA)
- Western blotting
- Dot- and slot-immunoblotting
- Immunohistochemistry

### CHARACTERIZATION

To ensure lot-to-lot consistency, each batch of product is tested by ELISA, FLISA, and/or flow cytometry for conformance to characteristics of a standard reference reagent.

### WORKING DILUTIONS

<b>Immunofluorescence</b>	FITC conjugate	≤ 1 µg/10 <sup>6</sup> cells
	R-PE conjugate	≤ 0.1 µg/10 <sup>6</sup> cells
	APC conjugate	≤ 0.1 µg/10 <sup>6</sup> cells
	R-PE-CY™7	≤ 0.1 µg/10 <sup>6</sup> cells
<b>ELISA:</b>	AP conjugate	1:2,000-1:4,000
	HRP conjugate	1:4,000-1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
<b>Other Applications:</b>	Since applications vary, you should determine the optimum working dilution of the product that is appropriate for your specific need.	

***For Research Use Only. Not for Diagnostic or Therapeutic Use.***

## HANDLING AND STORAGE

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 1.0 mg in 1.0 mL PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50mM Tris/1mM MgCl<sub>2</sub>/50% Glycerol, pH 8.0, containing 0.1% NaN<sub>3</sub> as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (R-PE) and allophycocyanin (APC) conjugates are supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The R-phycoerythrin-Cyanine 7 (R-PE-CY<sup>TM</sup>7) conjugate is supplied as 0.25 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## WARNING

Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.

\*Cy<sup>TM</sup>7 is a trademark of Amersham Biosciences Corp. Cy<sup>TM</sup>7 is for non-commercial research use only, not for therapeutic or in vivo applications. Other use needs license from Amersham Biosciences Corp., under U.S. Patent Nos. 4,981,977 and 5,268,486 and other patents pending. This material (or portions of this material) is subject to proprietary rights of Amersham Biosciences Corp. and Carnegie Mellon University and made and sold under license from Amersham Biosciences Corp.. This product is licensed for sale only for research. It is not licensed for any other use. There is no implied license hereunder for any commercial use. Commercial Use shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it. 2. sale, lease, license or other grant of rights to use this material or any material derived or produced from it. 3. Use of this material to perform services for a fee for third parties. If you require a commercial license to use this material and do not have one, return this material, unopened to Southern Biotechnology, 160A Oxmoor Blvd, Birmingham, AL 35209, USA and any money paid for the material will be refunded.