



Goat Anti-Rhesus IgG (H+L)

<u>Cat. No.</u>	<u>Form</u>	<u>Quantity</u>
6200-01	Purified (UNLB) Antibody	1.0 mg
6200-02	Fluorescein (FITC) Conjugate	1.0 mg
6200-03	Rhodamine (TRITC) Conjugate	1.0 mg
6200-04	Alkaline Phosphatase (AP) Conjugate	1.0 mL
6200-05	Horseradish Peroxidase (HRP) Conjugate	1.0 mL
6200-08	Biotin (BIOT) Conjugate	1.0 mg

DESCRIPTION

Source: Pooled antisera from goats hyperimmunized with rhesus IgG.
Cross Absorption: NA
Purification: Affinity chromatography on rhesus IgG covalently linked to agarose.
Specificity: Reacts with the heavy and light chains of rhesus IgG as demonstrated by ELISA. May also react with the light chains of other rhesus immunoglobulins, including IgM.

RESEARCH APPLICATIONS

- Immunofluorescent staining
- Enzyme-Linked-Immunosorbent-Assay (ELISA)
- Western blotting
- Dot- and slot-immunoblotting
- Immunohistochemistry

CHARACTERIZATION

To ensure lot-to-lot consistency, each batch of product is tested by ELISA and/or Fluorescent ELISA for conformance to characteristics of a standard reference reagent.

WORKING DILUTIONS

Immunofluorescence	Purified antibody, FITC, and TRITC conjugates	$\leq 1 \mu\text{g}/10^6$ cells
ELISA:	AP conjugate	1:2,000-1:4,000
	HRP conjugate	1:4,000-1:8,000
	BIOT conjugate	1:5,000-1:20,000

Other Applications: Since applications vary, you should determine the optimum working dilution of the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

HANDLING AND STORAGE

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) and rhodamine (TRITC) conjugates are supplied as 1.0 mg in 1.0 mL PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50mM Tris/1mM MgCl₂/50% Glycerol, pH 8.0, containing 0.1% NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

WARNING

Reagents contain sodium azide. Sodium azide is very toxic if ingested or inhaled. Avoid contact with skin, eyes, or clothing. Wear eye or face protection when handling. If skin or eye contact occurs, wash with copious amounts of water. If ingested or inhaled, contact a physician immediately. Sodium azide yields toxic hydrazoic acid under acidic conditions. Dilute azide-containing compounds in running water before discarding to avoid accumulation of potentially explosive deposits in lead or copper plumbing.