



## Protein G

Cat. No.	Form	Quantity
7406-01S	Purified (UNLB) Protein G	1.0 mg
7406-01L	Purified (UNLB) Protein G	5.0 mg
7406-05	Horseradish Peroxidase (HRP) Conjugate	1.0 mL

### DESCRIPTION

- Source** Recombinant, from *Streptococcus* Strain G148, produced in *E. coli*.
- Mr/Structure** 22 KD. Protein G is a monomeric protein lacking cystine residues; on SDS gel, it migrates to 32KD.
- Specificity** Protein G binds the Fc portion of IgGs from many species including all human, and mouse (IgG<sub>1</sub>, weakly) subclasses. It does not bind IgM, IgA, IgE or serum albumin.

### RESEARCH APPLICATIONS

- Affinity chromatography
- Immunoprecipitation
- Gamma immunoglobulin capture
- Conjugation on solid phase or others to detect gamma immunoglobulin
- Enzyme-Linked-Immunosorbent-Assay (ELISA)

### CHARACTERIZATION

To insure lot-to-lot consistency, each batch of product is tested by SDS-PAGE and/or ELISA to conform to the characteristics of a standard reference reagent.

### WORKING DILUTIONS

**ELISA** HRP conjugate 1:4,000-1:8,000

**Other Applications:** Since applications vary, you should determine the optimum working dilution of the product that is appropriate for your specific need.

### HANDLING AND STORAGE

- The purified Protein G is supplied as either 1.0 mg or 5.0 mg in lyophilized form. *No BSA, preservatives or amine-containing buffer salts have been added.* Store up to one year at -20°C; or after reconstitution in phosphate buffered saline to highest concentration possible, store up to one month at 2-8°C.
- Protein G-HRP conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. *No other preservatives have been added.* Store at 2-8°C or long-term at -20°C.
- Reagents are stable for the period shown on the label if stored as directed.

***For Research Use Only. Not for Diagnostic or Therapeutic Use.***

## REFERENCES

1. Saegerman, C., L. De Waele, D. Gilson, J. Godfroid, P. Thiange, P. Michel, B. Limbourg, T.K. Vo, J. Limet, J.J. Letesson, and D. Berkvens. 2004. *Vet. Microbiol.* 100:91.
2. Qi, Y., Z. Yan, and J. Huang. 2001. *J. Biochem. Biophys. Methods* 49:263.
3. Sugiura, T., H. Imahawa, and T. Kondo. 2000. *J. Chromatogr. B. Biomed. Sci. Appl.* 742:327.
4. Yan. Z., and J. Huang. 2000. *J. Immunol. Methods* 237:203.
5. Aybay, C., and T. Imir. 2000. *J. Immunol. Methods* 233:77.
6. Bastisa-Corcuera, F.D., K.H. Nielsen, and L.B. Corbeil. 1999. *Vet. Immunol. Immunopathol.* 71:143.
7. And, Q., G. Luo, J. Ou, and W.S. Yeung. 1999. *J. Chromatogr. A.* 848:139.
8. Goward CR, Murphy JP, Atkinson T, Barstow DA Expression and purification of a truncated recombinant streptococcal protein G. *Biochem J.* 1990 Apr 1;267(1):171-7.
9. Guss B, Eliasson M, Olsson A, Uhlén M, Frej AK, Jörnvall H, Flock JI, Lindberg M. Structure of the IgG-binding regions of streptococcal protein G. *EMBO J.* 1986 Jul;5(7):1567-75.