



Rat Anti-Human IL-13

Cat. No.	Format	Size
10125-01	Purified (UNLB)	0.5 mg
10125-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg

Overview

Clone	JES10-5A2
Isotype	Rat IgG _{1κ}
Immunogen	COS-expressed human IL-13
Specificity	Human IL-13
Alternate Name(s)	Interleukin-13, NC30

Applications

ELISA-Capture – Quality tested ^{2,3}
ELISA-Detection – Reported in literature ¹
ELISPOT-Detection – Reported in literature ⁴
FC – Reported in literature ⁵⁻¹⁰
IHC-FS – Reported in literature ¹¹⁻¹³
ICC – Reported in literature ¹⁴
Neut – Reported in literature ¹⁵

Note – May be paired with the biotinylated clone SB126d (SB Cat. No. 15930-08) in a sandwich ELISA

Working Dilutions

ELISA	Purified (UNLB) antibody	≤ 5 µg/mL
--------------	--------------------------	-----------

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: info@southernbiotech.com • Website: www.southernbiotech.com

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of PBS. Contains no preservative; handle under aseptic conditions. Store at 2-8°C or aliquot into smaller volumes and store at -20°C. Avoid multiple freeze / thaw cycles.
- Reagents are stable for the period shown on the label if stored as directed.

References

1. Mascart F, Verscheure V, Malfroot A, Hainaut M, Piérard D, Temerman S, et al. Bordetella pertussis infection in 2-month-old infants promotes type 1 T cell responses. *J Immunol.* 2003;170:1504-9. (ELISA-Detection)
2. Zhao W, Oskeritzian CA, Pozez AL, Schwartz LB. Cytokine production by skin-derived mast cells: endogenous proteases are responsible for degradation of cytokines. *J Immunol.* 2005;175:2635-42. (ELISA-Capture)
3. Janke M, Witsch EJ, Mages HW, Hutloff A, Kroczeck RA. Eminent role of ICOS costimulation for T cells interacting with plasmacytoid dendritic cells. *Immunology.* 2006;118:353-60. (ELISA-Capture)
4. Marotti B, Rocco A, De Colibus P, Compare D, de Nucci G, Staibano S, et al. Interleukin-13 mucosal production in Helicobacter pylori-related gastric diseases. *Dig Liver Dis.* 2008;40:240-7. (ELISPOT-Detection)
5. Gereda JE, Leung DY, Thatayatikom A, Streib JE, Price MR, Klinnert MD, et al. Relation between house-dust endotoxin exposure, type 1 T-cell development, and allergen sensitisation in infants at high risk of asthma. *Lancet.* 2000;355:1680-3. (FC)
6. Yamamoto J, Adachi Y, Onoue Y, Adachi YS, Okabe Y, Itazawa T, et al. Differential expression of the chemokine receptors by the Th1- and Th2-type effector populations within circulating CD4⁺ T cells. *J Leukoc Biol.* 2000;68:568-74. (FC)
7. Kim CH, Kunkel EJ, Boisvert J, Johnston B, Campbell JJ, Genovese MC, et al. Bonzo/CXCR6 expression defines type 1-polarized T-cell subsets with extralymphoid tissue homing potential. *J Clin Invest.* 2001;107:595-601. (FC)
8. Iwasaki M, Nagata K, Takano S, Takahashi K, Ishii N, Ikezawa Z. Association of a new-type prostaglandin D2 receptor CRTH2 with circulating T helper 2 cells in patients with atopic dermatitis. *J Invest Dermatol.* 2002;119:609-16. (FC)
9. Annunziato F, Cosmi L, Liotta F, Lazzeri E, Manetti R, Vanini V, et al. Phenotype, localization, and mechanism of suppression of CD4⁺CD25⁺ human thymocytes. *J Exp Med.* 2002;196:379-87. (FC)
10. Cousins DJ, Lee TH, Staynov DZ. Cytokine coexpression during human Th1/Th2 cell differentiation: direct evidence for coordinated expression of Th2 cytokines. *J Immunol.* 2002;169:2498-506. (FC)
11. Ulfgrén A, Lindblad S, Klareskog L, Andersson J, Andersson U. Detection of cytokine producing cells in the synovial membrane from patients with rheumatoid arthritis. *Ann Rheum Dis.* 1995;54:654-61. (IHC-FS)
12. Sparrelid E, Emanuel D, Fehniger T, Andersson U, Andersson J. Interstitial pneumonitis in bone marrow transplant recipients is associated with local production of TH2-type cytokines and lack of T cell-mediated cytotoxicity. *Transplantation.* 1997;63:1782-9. (IHC-FS)
13. Knörr C, Amrehn C, Seeberger H, Rosenwald A, Stilgenbauer S, Ott G, et al. Expression of costimulatory molecules in low-grade mucosa-associated lymphoid tissue-type lymphomas in vivo. *Am J Pathol.* 1999;155:2019-27. (IHC-FS)
14. Dakhama A, Collins ML, Ohnishi H, Goleva E, Leung DY, Alam R, et al. IL-13-producing BLT1-positive CD8 cells are increased in asthma and are associated with airway obstruction. *Allergy.* 2013;68:666-73. (ICC)
15. Punnonen J, Yssel H, de Vries JE. The relative contribution of IL-4 and IL-13 to human IgE synthesis induced by activated CD4⁺ or CD8⁺ T cells. *J Allergy Clin Immunol.* 1997;100:792-801. (Neut)