




Goat Anti-Rabbit Ig, Human ads

Cat. No.	Format	Size
4010-01	Purified (UNLB)	1.0 mg
4010-02	Fluorescein (FITC)	1.0 mg
4010-03	Rhodamine (TRITC)	1.0 mg
4010-04	Alkaline Phosphatase (AP)	1.0 mL
4010-05	Horseradish Peroxidase (HRP)	1.0 mL
4010-06	β -galactosidase (BGAL)	1.0 mL
4010-07	Texas Red [®] (TXRD)	1.0 mg
4010-08	Biotin (BIOT)	1.0 mg
4010-09	R-phycoerythrin (PE)	0.5 mg
4010-09S	R-phycoerythrin (PE)	0.25 mg
4010-30	Alexa Fluor [®] 488 (AF488)	1.0 mg
4010-31	Alexa Fluor [®] 647 (AF647)	1.0 mg
4010-32	Alexa Fluor [®] 555 (AF555)	1.0 mg



ELISA plate was coated with purified rabbit IgG and IgM and human IgG, IgM, and IgA. Immunoglobulins were detected with serially diluted Goat Anti-Rabbit Ig, Human ads-BIOT (SB Cat. No. 4010-08) followed by Streptavidin-HRP (SB Cat. No. 7100-05).

Description

Specificity	Reacts with the heavy and light chains of rabbit IgG and IgM
Source	Pooled antisera from goats hyperimmunized with rabbit IgG
Cross Adsorption	Human immunoglobulins and pooled sera; may react with immunoglobulins from other species
Purification	Affinity chromatography on pooled rabbit IgG covalently linked to agarose

Applications

Quality tested applications include –

ELISA ¹⁻³
FLISA
FC ⁴

Other referenced applications include –

IHC-FS ⁵⁻⁷
IHC-PS ^{3,8}
ICC ⁹⁻¹⁴
WB ¹⁵⁻²⁵
IP ²⁶

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BGAL conjugate	1:500 – 1:2,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE and AF647 conjugates	≤ 1 µg/mL
Flow Cytometry	FITC and BIOT conjugates	≤ 1 µg/10 ⁶ cells
	PE and AF647 conjugates	≤ 0.1 µg/10 ⁶ cells
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL	
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The β-galactosidase (BGAL) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.5 mg in 1.0 mL or 0.25 mg in 0.5 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Fotopoulou S, Oikonomou N, Grigorieva E, Nikitopoulou I, Paparountas T, Thanassopoulou A, et al. ATX expression and LPA signalling are vital for the development of the nervous system. *Dev Biol.* 2010;339:451-64. (ELISA)
2. Jain S, Singh SR, Horn DW, Davis VA, Ram MK, Pillai S. Development of an antibody functionalized carbon nanotube biosensor for foodborne bacterial pathogens. *J Biosens Bioelectron.* 2012;S11:002. (ELISA)
3. Béteemps D, Verchère J, Brot S, Morignat E, Bousset L, Gaillard D, et al. Alpha-synuclein spreading in M83 mice brain revealed by detection of pathological α-synuclein by enhanced ELISA. *Acta Neuropathol Commun.* 2014;2:29. (ELISA, IHC-PS)
4. Hautefort I, Proença MJ, Hinton JC. Single-copy green fluorescent protein gene fusions allow accurate measurement of Salmonella gene expression in vitro and during infection of mammalian cells. *Appl Environ Microbiol.* 2003;69:7480-91. (FC)
5. Perry LL, Feilzer K, Portis JL, Caldwell HD. Distinct homing pathways direct T lymphocytes to the genital and intestinal mucosae in Chlamydia-infected mice. *J Immunol.* 1998;160:2905-14. (IHC-FS)
6. Manley HA, Lennon VA. Endoplasmic reticulum membrane-sorting protein of lymphocytes (BAP31) is highly expressed in neurons and discrete endocrine cells. *J Histochem Cytochem.* 2001;49:1235-43. (IHC-FS)
7. Gruver AL, Ventevogel MS, Sempowski GD. Leptin receptor is expressed in thymus medulla and leptin protects against thymic remodeling during endotoxemia-induced thymus involution. *J Endocrinol.* 2009;203:75-85. (IHC-FS)
8. Sang Y, Zhou T, Li H, Ran Y, Jiang D, Zheng F, et al. A new long-acting GLP-1 derivative KTP ameliorates hyperglycemia and dyslipidemia and improves pancreas and fatty liver in db/db mice. *Chin Sci Bull.* 2013;58:2447-53. (IHC-PS)
9. Campana D, van Dongen JJ, Mehta A, Coustan-Smith E, Wolvers-Tettero IL, Ganeshaguru K, et al. Stages of T-cell receptor protein expression in T-cell acute lymphoblastic leukemia. *Blood.* 1991;77:1546-54. (ICC)
10. Ivan M, Bond JA, Prat M, Comoglio PM, Wynford-Thomas D. Activated ras and ret oncogenes induce over-expression of c-met (hepatocyte growth factor receptor) in human thyroid epithelial cells. *Oncogene.* 1997;14:2417-23. (ICC)
11. Alonso M, Tamasdan C, Miller DC, Newcomb EW. Flavopiridol induces apoptosis in glioma cell lines independent of retinoblastoma and p53 tumor suppressor pathway alterations by a caspase-independent pathway. *Mol Cancer Ther.* 2003;2:139-50. (ICC)
12. Lechuga CG, Hernández-Nazara ZH, Hernández E, Bustamante M, Desierto G, Cotty A, et al. PI3K is involved in PDGF-β receptor upregulation post-PDGF-BB treatment in mouse HSC. *Am J Physiol Gastrointest Liver Physiol.* 2006;291:G1051-61. (ICC)
13. Haegens A, Vernooy JH, Heeringa P, Mossman BT, Wouters EF. Myeloperoxidase modulates lung epithelial responses to pro-inflammatory agents. *Eur Respir J.* 2008;31:252-60. (ICC)
14. Bishop RK, Valle Osegura CA, Spencer JV. Human Cytomegalovirus interleukin-10 promotes proliferation and migration of MCF-7 breast cancer cells. *Cancer Cell Microenviron.* 2015;2(1):pii: e678. (ICC)
15. Webb CF, Yamashita Y, Ayers N, Evetts S, Paulin Y, Conley ME, et al. The transcription factor Bright associates with Bruton's tyrosine kinase, the defective protein in immunodeficiency disease. *J Immunol.* 2000;165:6956-65. (WB)
16. Pilling D, Buckley CD, Salmon M, Gomer RH. Inhibition of fibrocyte differentiation by serum amyloid P. *J Immunol.* 2003;171:5537-46. (WB)
17. Bai Q, Garver JA, Hukriede NA, Burton EA. Generation of a transgenic zebrafish model of Tauopathy using a novel promoter element derived from the zebrafish eno2 gene. *Nucleic Acids Res.* 2007;35:6501-16. (WB)
18. Hangen E, De Zio D, Bordi M, Zhu C, Dessen P, Caffin F, et al. A brain-specific isoform of mitochondrial apoptosis-inducing factor: AIF2. *Cell Death Differ.* 2010;17:1155-66. (WB)
19. Palatinus JA, O'Quinn MP, Barker RJ, Harris BS, Jourdan J, Gourdie RG. ZO-1 determines adherens and gap junction localization at intercalated disks. *Am J Physiol Heart Circ Physiol.* 2011;300:H583-94. (WB)
20. Sommer A, Christensen E, Schwenger S, Seul R, Haas D, Olbrich H, et al. The molecular basis of aminoacylase 1 deficiency. *Biochim Biophys Acta.* 2011;1812:685-90. (WB)
21. Babcock AS, Anderson AE, Rice CD. Indirubin-3'-(2,3 dihydroxypropyl)-oximether (E804) is a potent modulator of LPS-stimulated macrophage functions. *Toxicol Appl Pharmacol.* 2013;266:157-66. (WB)
22. Chen Z, Jolley B, Caldwell C, Gallie DR. Eukaryotic translation initiation factor eIFiso4G is required to regulate violaxanthin de-epoxidase expression in Arabidopsis. *J Biol Chem.* 2014;289:13926-36. (WB)
23. Andersen TR, Schmidt JF, Thomassen M, Hornstrup T, Frandsen U, Randers MB, et al. A preliminary study: effects of football training on glucose control, body composition, and performance in men with type 2 diabetes. *Scand J Med Sci Sports.* 2014;24 Suppl 1:43-56. (WB)
24. Palatinus JA, Gourdie RG. Diabetes increases cryoinjury size with associated effects on Cx43 gap junction function and phosphorylation in the mouse heart. *J Diabetes Res.* 2015;2015:510935. (WB)
25. Xayarath B, Alonzo F 3rd, Freitag NE. Identification of a peptide-pheromone that enhances Listeria monocytogenes escape from host cell vacuoles. *PLoS Pathog.* 2015;11(3):e1004707. (WB)
26. Jurivich DA, Pachetti C, Qiu L, Welk JF. Salicylate triggers heat shock factor differently than heat. *J Biol Chem.* 1995;270:24489-95. (IP)

Texas Red® is a registered trademark of Molecular Probes, Inc.
Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

TB4010
25-Aug-21

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • **Fax:** 205.945.8768

Email: info@southernbiotech.com • **Website:** www.southernbiotech.com