
SouthernBiotech



Goat Anti-Canine IgG(H+L)

Cat. No.	Format	Size
6070-01	Purified (UNLB)	1.0 mg
6070-02	Fluorescein (FITC)	1.0 mg
6070-04	Alkaline Phosphatase (AP)	1.0 mL
6070-05	Horseradish Peroxidase (HRP)	1.0 mL
6070-08	Biotin (BIOT)	1.0 mg
6070-30	Alexa Fluor® 488 (AF488)	1.0 mg



ELISA plate was coated with purified canine IgG. Immunoglobulin was detected with Goat Anti-Canine IgG(H+L)-AP (SB Cat. No. 6070-04).

Description

Specificity Reacts with the heavy and light chains of canine IgG

Source Pooled antisera from goats hyperimmunized with canine IgG

Cross Adsorption None; may react with immunoglobulins from other species and the light chains of other canine

immunoglobulins

Purification Affinity chromatography on canine IgG covalently linked to agarose

Applications

Quality tested applications include -

ELISA 1,2 FLISA

Other referenced applications include -

IHC-FS ³ ICC ⁴ WB ⁵⁻⁸

Working Dilutions

ELISA	AP conjugate	1:2.000 – 1:4.000

HRP conjugate 1:4,000 – 1:8,000 BIOT conjugate 1:5,000 – 1:20,000

FLISA FITC and AF488 conjugates 1:200 – 1:400

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is

appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added. Store at 2-8°C.
- The fluorescein (FITC) and Alexa Fluor® 488 (AF488) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

- 1. Kakkis E, Lester T, Yang R, Tanaka C, Anand V, Lemontt J, et al. Successful induction of immune tolerance to enzyme replacement therapy in canine mucopolysaccharidosis I. Proc Natl Acad Sci USA. 2004;101:829-34. (ELISA)
- 2. Dickson P, Peinovich M, McEntee M, Lester T, Le S, Krieger A, et al. Immune tolerance improves the efficacy of enzyme replacement therapy in canine mucopolysaccharidosis I. J Clin Invest. 2008;118:2868-76. (ELISA)
- 3. Wiberg ME, Saari SA, Westermarck E, Meri S. Cellular and humoral immune responses in atrophic lymphocytic pancreatitis in German shepherd dogs and rough-coated collies. Vet Immunol Immunopathol. 2000;76:103-15. (IHC-FS)
- Ulchar I, Celeska I, Stefanovska J, Jakimovska A. Hematological and biochemical parameters in symptomatic and asymptomatic leishmaniaseropositive dogs. Mac Vet Rev. 2015;38:i-viii. (ICC)
- Mathis A, Åkerstedt J, Tharaldsen J, Ødegaard Ø, Deplazes P. Isolates of Encephalitozoon cuniculi from farmed blue foxes (Alopex Iagopus) from Norway differ from isolates from Swiss domestic rabbits (Oryctolagus cuniculus). Parasitol Res. 1996;82:727-30. (WB)
- Rasmussen UB, Benchaibi M, Meyer V, Schlesinger Y, Schughart K. Novel human gene transfer vectors: evaluation of wild-type and recombinant animal adenoviruses in human-derived cells. Hum Gene Ther. 1999;10:2587-99. (WB)
- 7. Müller-Doblies UU, Herzog K, Tanner I, Mathis A, Deplazes P. First isolation and characterisation of Encephalitozoon cuniculi from a free-ranging rat (Rattus norvegicus). Vet Parasitol. 2002;107:279-85. (WB)
- 8. Burk SV. Detection of antibodies against Parascaris equorum excretory-secretory antigens [dissertation]. Lexington (USA): University of Kentucky, 2013 (WB)

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

Email: info@southernbiotech.com • Website: www.southernbiotech.com