




Rat Anti-Human IL-10

Cat. No.	Format	Size
10110-01	Purified (UNLB)	0.5 mg
10110-08	Biotin (BIOT)	0.5 mg



Standard curve generated with Rat Anti-Human IL-10-UNLB (SB Cat. No. 10100-01; Clone JES3-9D7) and Rat Anti-Human IL-10-BIOT (SB Cat. No. 10110-08; Clone JES3-12G8) followed by Streptavidin-HRP (SB Cat. No. 7100-05)

Overview

Clone	JES3-12G8
Isotype	Rat IgG _{2a} κ
Immunogen	COS-expressed human IL-10
Specificity	Human/Viral/Cynomolgus IL-10
Alternate Name(s)	Interleukin-10, cytokine synthesis inhibitory factor, CSIF, TGIF, B-TCGF

Applications

ELISA-Detection – Quality tested^{1-4,6}
 ELISPOT-Detection – Reported in literature⁵⁻⁷
 IHC-FS – Reported in literature⁸⁻¹⁰
 IHC-PS – Reported in literature¹¹
 WB – Reported in literature^{4,12}
 Neut – Reported in literature^{1,12,13}
 Multiplex-Detection – Reported in literature¹⁴

Note – May be paired with the purified clone JES3-9D7 (SB Cat. No. 10100-01) in a sandwich ELISA

Working Dilutions

ELISA	BIOT conjugate	1:2,000 – 1:4,000
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg labeled antibody in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

References

1. Abrams JS, Roncarolo M, Yssel H, Andersson U, Gleich GJ, Silver JE. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev.* 1992;127:5-24. (ELISA-Detection, Neut)
2. Gottlieb WH, Abrams JS, Watson JM, Velu TJ, Berek JS, Martínez-Maza O. Presence of interleukin 10 (IL-10) in the ascites of patients with ovarian and other intra-abdominal cancers. *Cytokine.* 1992;4:385-90. (ELISA-Detection, Viral Reactivity)
3. Abrams JS. Immunoenzymetric assay of mouse and human cytokines using NIP-labeled anti-cytokine antibodies. *Curr Protoc Immunol.* 2001;6.20:1-15. (ELISA-Detection)
4. Kawamoto S, Nitta Y, Tashiro F, Nakano A, Yamato E, Tahara H, et al. Suppression of T_h1 cell activation and prevention of autoimmune diabetes in NOD mice by local expression of viral IL-10. *Int Immunol.* 2001;13:685-94. (ELISA-Detection, WB, Viral Reactivity)
5. Hansson T, Dannæus A, Klareskog L. Cytokine-producing cells in peripheral blood of children with coeliac disease secrete cytokines with a type 1 profile. *Clin Exp Immunol.* 1999;116:246-50. (ELISPOT-Detection)
6. Guerkov RE, Targoni OS, Kreher CR, Boehm BO, Herrera MT, Tary-Lehmann M, et al. Detection of low-frequency antigen-specific IL-10-producing CD4⁺ T cells via ELISPOT in PBMC: cognate vs. nonspecific production of the cytokine. *J Immunol Methods.* 2003;279:111-21. (ELISA-Detection, ELISPOT-Detection)
7. Klinman D. ELISPOT assay to detect cytokine-secreting murine and human cells. *Curr Protoc Immunol.* 2008;6.19:1-9. (ELISPOT-Detection)
8. Andersson J, Abrams J, Björk L, Funa K, Litton M, Ågren K, et al. Concomitant in vivo production of 19 different cytokines in human tonsils. *Immunology.* 1994;83:16-24. (IHC-FS)
9. Ulfgrén A, Lindblad S, Klareskog L, Andersson J, Andersson U. Detection of cytokine producing cells in the synovial membrane from patients with rheumatoid arthritis. *Ann Rheum Dis.* 1995;54:654-61. (IHC-FS)
10. Bergeron A, Soler P, Kambouchner M, Loiseau P, Milleron B, Valeyre D, et al. Cytokine profiles in idiopathic pulmonary fibrosis suggest an important role for TGF- β and IL-10. *Eur Respir J.* 2003;22:69-76. (IHC-FS)
11. Bellone G, Carbone A, Smirne C, Scirelli T, Buffolino A, Novarino A, et al. Cooperative induction of a tolerogenic dendritic cell phenotype by cytokines secreted by pancreatic carcinoma cells. *J Immunol.* 2006;177:3448-60. (IHC-PS)
12. Brodeur ND, Spencer JV. Antibodies to human IL-10 neutralize ebvIL-10-mediated cytokine suppression but have no effect on cmvIL-10 activity. *Virus Res.* 2010;153:265-8. (WB, Neut, Viral Reactivity)
13. Groux H, O'Garra A, Bigler M, Rouleau M, Antonenko S, de Vries JE, et al. A CD4⁺ T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature.* 1997;389:737-42. (Neut)
14. Hutchinson KL, Villinger F, Miranda ME, Ksiazek TG, Peters CJ, Rollin PE. Multiplex analysis of cytokines in the blood of cynomolgus macaques naturally infected with Ebola virus (Reston serotype). *J Med Virol.* 2001;65:561-6. (Multiplex-Detection, Cynomolgus Reactivity)