




## Goat Anti-Mouse IgG, Human ads

Cat. No.	Format	Size
1030-01	Purified (UNLB)	1.0 mg
1030-02	Fluorescein (FITC)	1.0 mg
1030-03	Rhodamine (TRITC)	1.0 mg
1030-04	Alkaline Phosphatase (AP)	1.0 mL
1030-05	Horseradish Peroxidase (HRP)	1.0 mL
1030-06	$\beta$ -galactosidase (BGAL)	1.0 mL
1030-07	Texas Red <sup>®</sup> (TXRD)	1.0 mg
1030-08	Biotin (BIOT)	1.0 mg
1030-09	R-phycoerythrin (PE)	0.5 mg
1030-09S	R-phycoerythrin (PE)	0.25 mg
1030-15	Cyanine 5 (CY5)	1.0 mg
1030-30	Alexa Fluor <sup>®</sup> 488 (AF488)	1.0 mg
1030-31	Alexa Fluor <sup>®</sup> 647 (AF647)	1.0 mg
1030-32	Alexa Fluor <sup>®</sup> 555 (AF555)	1.0 mg



ELISA plate was coated with purified mouse IgG, IgM, and IgA. Immunoglobulins were detected with serially diluted Goat Anti-Mouse IgG, Human ads-HRP (SB Cat. No. 1030-05).

### Description

<b>Specificity</b>	Reacts with the heavy chains of mouse IgG <sub>1</sub> , IgG <sub>2a</sub> , IgG <sub>2b</sub> , IgG <sub>2c</sub> , and IgG <sub>3</sub>
<b>Source</b>	Pooled antisera from goats hyperimmunized with mouse IgG
<b>Cross Adsorption</b>	Mouse IgM and IgA; human immunoglobulins and pooled sera; may react with immunoglobulins from other species
<b>Purification</b>	Affinity chromatography on mouse IgG covalently linked to agarose

### Applications

Quality tested applications include –

ELISA<sup>1-10</sup>  
 FLISA  
 FC<sup>5,11-13</sup>

Other referenced applications include –

ELISPOT<sup>1,3-6</sup>  
 IHC-FS<sup>2,14,15</sup>  
 IHC-PS<sup>16-18</sup>  
 ICC<sup>19-22</sup>  
 WB<sup>7,21,23-25</sup>  
 Multiplex<sup>26</sup>

### Working Dilutions

<b>ELISA</b>	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BGAL conjugate	1:500
	BIOT conjugate	1:5,000 – 1:20,000
<b>FLISA</b>	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE, CY5, and AF647 conjugates	≤ 1 µg/mL
<b>Flow Cytometry</b>	FITC, BIOT, and AF488 conjugates	≤ 1 µg/10 <sup>6</sup> cells
	PE, CY5, and AF647 conjugates	≤ 0.1 µg/10 <sup>6</sup> cells
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL	
<b>Other Applications</b>	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

## Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added. Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Cyanine 5 (CY5), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50 mM Tris/1 mM MgCl<sub>2</sub>/50% glycerol, pH 8.0, containing NaN<sub>3</sub> as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The β-galactosidase (BGAL) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS containing NaN<sub>3</sub> as preservative. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.5 mg in 1.0 mL or 0.25 mg in 0.5 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

## References

1. van Ginkel FW, Wahl SM, Kearney JF, Kweon M, Fujihashi K, Burrows PD, et al. Partial IgA-deficiency with increased Th2-type cytokines in TGF-β1 knockout mice. *J Immunol.* 1999;163:1951-7. (ELISA, ELISPOT)
2. Dunn LA, Upcroft JA, Fowler EV, Matthews BS, Upcroft P. Orally administered *Giardia duodenalis* extracts enhance an antigen-specific antibody response. *Infect Immun.* 2001;69:6503-10. (ELISA, IHC-FS)
3. Lagrota-Candido J, Vasconcellos R, Cavalcanti M, Bozza M, Savino W, Quirico-Santos T. Resolution of skeletal muscle inflammation in mdx dystrophic mouse is accompanied by increased immunoglobulin and interferon-γ production. *Int J Exp Path.* 2002;83:121-32. (ELISA, ELISPOT)
4. Hovden A, Cox RJ, Haaheim LR. Whole influenza virus vaccine is more immunogenic than split influenza virus vaccine and induces primarily an IgG2a response in BALB/c mice. *Scand J Immunol.* 2005;62:36-44. (ELISA, ELISPOT)
5. Conway KL, Kuballa P, Khor B, Zhang M, Shi HN, Virgin HW, et al. ATG5 regulates plasma cell differentiation. *Autophagy.* 2013;9:528-37. (ELISA, ELISPOT, FC)
6. Koutsouanos DG, Esser ES, McMaster SR, Kalluri P, Lee J, Prausnitz MR, et al. Enhanced immune responses by skin vaccination with influenza subunit vaccine in young hosts. *Vaccine.* 2015;33:4675-82. (ELISA, ELISPOT)
7. Yolken RH, Jones-Brando L, Dunigan DD, Kannan G, Dickerson F, Severance E, et al. Chlorovirus ATCV-1 is part of the human oropharyngeal virome and is associated with changes in cognitive functions in humans and mice. *Proc Natl Acad Sci USA.* 2014;111:16106-11. (ELISA, WB)
8. Zhou Y, Maharaj PD, Mallajosyula JK, McCormick AA, Kearney CM. In planta production of flock house virus transencapsidated RNA and its potential use as a vaccine. *Mol Biotechnol.* 2015;57:325-36. (ELISA)
9. Zhao H, Bauzon F, Bi E, Yu JJ, Fu H, Lu Z, et al. Substituting threonine 187 with alanine in p27Kip1 prevents pituitary tumorigenesis by two-hit loss of Rb1 and enhances humoral immunity in old age. *J Biol Chem.* 2015;290:5797-809. (ELISA)
10. Schubert RD, Hu Y, Kumar G, Szeto S, Abraham P, Winderl J, et al. IFN-β treatment requires B cells for efficacy in neuroautoimmunity. *J Immunol.* 2015;194:2110-6. (ELISA)
11. Chen J, Yin H, Xu J, Wang Q, Edelblum KL, Sciammas R, et al. Reversing endogenous alloreactive B cell GC responses with anti-CD154 or CTLA-4lg. *Am J Transplant.* 2013;13:2280-92. (FC)
12. Maxwell P, Melendez-Rodríguez F, Matchett KB, Aragonés J, Ben-Califa N, Jaekel H, et al. Novel antibodies directed against the human erythropoietin receptor: creating a basis for clinical implementation. *Br J Haematol.* 2015;168:429-42. (FC)
13. Brady AM, Spencer BL, Falsey AR, Nahm MH. Blood collection tubes influence serum ficolin-1 and ficolin-2 levels. *Clin Vaccine Immunol.* 2014;21:51-5. (FC)
14. Csencsits K, Burrell BE, Lu G, Eichwald EJ, Stahl GL, Bishop DK. The classical complement pathway in transplantation: unanticipated protective effects of C1q and role in inductive antibody therapy. *Am J Transplant.* 2008;8:1622-30. (IHC-FS)
15. Sic H, Kraus H, Madl J, Flittner K, von Münchow AL, Pieper K, et al. Sphingosine-1-phosphate receptors control B-cell migration through signaling components associated with primary immunodeficiencies, chronic lymphocytic leukemia, and multiple sclerosis. *J Allergy Clin Immunol.* 2014;134:420-8. (IHC-FS)
16. Zheng P, Zhao YX, Zhang AD, Kang C, Chen HC, Jin ML. Pathologic analysis of the brain from *Streptococcus suis* type 2 experimentally infected pigs. *Vet Pathol.* 2009;46:531-5. (IHC-PS)
17. McPhee CG, Sproule TJ, Shin D, Bubier JA, Schott WH, Steinbuck MP, et al. MHC class I family proteins retard systemic lupus erythematosus autoimmunity and B cell lymphomagenesis. *J Immunol.* 2011;187:4695-704. (IHC-PS)
18. Vujasinovic T, Pribic J, Kanjer K, Milosevic NT, Tomasevic Z, Milovanovic Z, et al. Gray-level co-occurrence matrix texture analysis of breast tumor images in prognosis of distant metastasis risk. *Microsc Microanal.* 2015;21:646-54. (IHC-PS)
19. Sjöberg M, Garoff H. Interactions between the transmembrane segments of the alphavirus E1 and E2 proteins play a role in virus budding and fusion. *J Virol.* 2003;77:3441-50. (ICC)
20. Roberts MS, Woods AJ, Dale TC, van der Sluijs P, Norman JC. Protein kinase B/Akt acts via glycogen synthase kinase 3 to regulate recycling of αvβ3 and α5β1 integrins. *Mol Cell Biol.* 2004;24:1505-15. (ICC)
21. Ma Y, Yates J, Liang Y, Lemon SM, Yi M. NS3 helicase domains involved in infectious intracellular hepatitis C virus particle assembly. *J Virol.* 2008;82:7624-39. (ICC, WB)
22. Mia MM, Bank RA. Paracrine factors of human amniotic fluid-derived mesenchymal stem cells show strong anti-fibrotic properties by inhibiting myofibroblast differentiation and collagen synthesis. *J Stem Cell Res Ther.* 2015;5:5. (ICC)
23. Friedman J, Alam SM, Shen X, Xia S, Stewart S, Anasti K, et al. Isolation of HIV-1-neutralizing mucosal monoclonal antibodies from human colostrum. *PLoS One.* 2012;7(5):e37648. (WB)
24. Bender BJ, Coen DM, Strang BL. Dynamic and nucleolin-dependent localization of human cytomegalovirus UL84 to the periphery of viral replication compartments and nucleoli. *J Virol.* 2014;88:11738-47. (WB)
25. Hethorn WR, Ciarlone SL, Filonova I, Rogers JT, Aguirre D, Ramirez RA, et al. Reelin supplementation recovers synaptic plasticity and cognitive deficits in a mouse model for Angelman syndrome. *Eur J Neurosci.* 2015;41:1372-80. (WB)
26. Hovden A, Brokstad KA, Major D, Wood J, Haaheim LR, Cox RJ. A pilot study of the immune response to whole inactivated avian influenza H7N1 virus vaccine in mice. *Influenza Other Respir Viruses.* 2009;3:21-8. (Multiplex)

Texas Red® is a registered trademark of Molecular Probes, Inc.

Cy® is a registered trademark of GE Healthcare.

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or [outlicensing@lifetech.com](mailto:outlicensing@lifetech.com).

TB1030

16-Nov-16

**Corporate Offices:** 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

**Tel:** 205.945.1774 • U.S. and Canada: 800.722.2255 • **Fax:** 205.945.8768

**Email:** [info@southernbiotech.com](mailto:info@southernbiotech.com) • **Website:** [www.southernbiotech.com](http://www.southernbiotech.com)