




Goat Anti-Mouse IgG_{2a}, Human ads

Cat. No.	Format	Size
1080-01	Purified (UNLB)	1.0 mg
1080-02	Fluorescein (FITC)	1.0 mg
1080-03	Rhodamine (TRITC)	1.0 mg
1080-04	Alkaline Phosphatase (AP)	1.0 mL
1080-05	Horseradish Peroxidase (HRP)	1.0 mL
1080-07	Texas Red [®] (TXRD)	1.0 mg
1080-08	Biotin (BIOT)	1.0 mg
1080-09	R-phycoerythrin (PE)	0.5 mg
1080-09S	R-phycoerythrin (PE)	0.25 mg
1080-10	R-phycoerythrin-Texas Red [®] (PE/TXRD)	0.25 mg
1080-11L	Allophycocyanin (APC)	0.5 mg
1080-11S	Allophycocyanin (APC)	0.25 mg
1080-13	Spectral Red [®] (SPRD)	0.25 mg
1080-15	Cyanine 5 (CY5)	1.0 mg
1080-17	R-phycoerythrin-Cyanine 7 (PE/CY7)	0.25 mg
1080-19	Allophycocyanin-Cyanine 7 (APC/CY7)	0.25 mg
1080-30	Alexa Fluor [®] 488 (AF488)	1.0 mg
1080-31	Alexa Fluor [®] 647 (AF647)	1.0 mg
1080-32	Alexa Fluor [®] 555 (AF555)	1.0 mg



FLISA plate was coated with purified mouse IgG₁, IgG_{2a}, IgG_{2b}, IgG₃, IgM, and IgA. Immunoglobulins were detected with serially diluted Goat Anti-Mouse IgG_{2a}, Human ads-APC (SB Cat. No. 1080-11).

Description

Specificity	Reacts with the heavy chain of mouse IgG _{2a}
Source	Pooled antisera from goats hyperimmunized with mouse IgG _{2a}
Cross Adsorption	Mouse IgG ₁ , IgG _{2b} , IgG ₃ , IgM, and IgA; human immunoglobulins and pooled sera; may react with immunoglobulins from other species
Purification	Affinity chromatography on mouse IgG _{2a} covalently linked to agarose

Applications

Quality tested applications include –

ELISA¹⁻⁹
 FLISA
 FC¹¹⁻¹³

Other referenced applications include –

ELISPOT^{3,10}
 IHC-FS^{2,14,15}
 IHC-PS^{16,17}
 ICC^{4,18-20}
 EM¹⁴
 WB^{1,21-23}
 SPR²⁴
 Purification²⁵

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE, APC, CY5, and AF647 conjugates	≤ 1 µg/mL
Flow Cytometry	FITC, BIOT, and AF488 conjugates	≤ 1 µg/10 ⁶ cells
	PE, PE/TXRD, APC, CY5, PE/CY7, APC/CY7, and AF647 conjugates	≤ 0.1 µg/10 ⁶ cells

For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Cyanine 5 (CY5), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. *No preservative added.* Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) and allophycocyanin (APC) conjugates are supplied as 0.5 mg in 1.0 mL or 0.25 mg in 0.5 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral Red® (SPRD), R-phycoerythrin-Texas Red® (PE/TXRD), R-phycoerythrin-Cyanine 7 (PE/CY7), and allophycocyanin-Cyanine 7 (APC/CY7) conjugates are supplied as 0.25 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Martin RM, Brady JL, Lew AM. The need for IgG2c specific antiserum when isotyping antibodies from C57BL/6 and NOD mice. *J Immunol Methods.* 1998;212:187-92. (ELISA, WB)
2. Richards HB, Satoh M, Jennette JC, Croker BP, Yoshida H, Reeves WH. Interferon- γ is required for lupus nephritis in mice treated with the hydrocarbon oil pristane. *Kidney Int.* 2001;60:2173-80. (ELISA, IHC-FS)
3. Cox RJ, Mykkeltvedt E, Robertson J, Haaheim LR. Non-lethal viral challenge of influenza haemagglutinin and nucleoprotein DNA vaccinated mice results in reduced viral replication. *Scand J Immunol.* 2002;55:14-23. (ELISA, ELISPOT)
4. Swiecki MK, Lisanby MW, Shu F, Turnbough CL Jr, Kearney JF. Monoclonal antibodies for Bacillus anthracis spore detection and functional analyses of spore germination and outgrowth. *J Immunol.* 2006;176:6076-84. (ELISA, ICC)
5. Arnaboldi PM, Roth-Walter F, Mayer L. Suppression of Th1 and Th17, but not Th2, responses in a CD8⁺ T cell mediated model of oral tolerance. *Mucosal Immunol.* 2009;2:427-38. (ELISA)
6. Faustino L, Fonseca DM, Florsheim EB, Resende RR, Lepique AP, Faquim-Mauro E, et al. Tumor necrosis factor-related apoptosis-inducing ligand mediates the resolution of allergic airway inflammation induced by chronic allergen inhalation. *Mucosal Immunol.* 2014;7:1199-208. (ELISA)
7. Hartung E, Becker M, Bachem A, Reeg N, Jäkel A, Hutloff A, et al. Induction of potent CD8 T cell cytotoxicity by specific targeting of antigen to cross-presenting dendritic cells in vivo via murine or human XCR1. *J Immunol.* 2015;194:1069-79. (ELISA)
8. Dabral N, Jain-Gupta N, Selem MN, Siranganathan N, Vemulapalli R. Overexpression of Brucella putative glycosyltransferase WbKa in B. abortus RB51 leads to production of exopolysaccharide. *Front Cell Infect Microbiol.* 2015;5:54. (ELISA)
9. Dabaghian M, Latifi AM, Tebianian M, Dabaghian F, Ebrahimi SM. A truncated C-terminal fragment of Mycobacterium tuberculosis HSP70 enhances cell-mediated immune response and longevity of the total IgG to influenza A virus M2e protein in mice. *Antiviral Res.* 2015;120:23-31. (ELISA)
10. Teichmann LL, Schenten D, Medzhitov R, Kashgarian M, Shlomchik MJ. Signals via the adaptor MyD88 in B cells and DCs make distinct and synergistic contributions to immune activation and tissue damage in lupus. *Immunity.* 2013;38:528-40. (ELISPOT)
11. Davis ME, Maxwell CV, Erf GF, Brown DC, Wistuba TJ. Dietary supplementation with phosphorylated mannans improves growth response and modulates immune function of weanling pigs. *J Anim Sci.* 2004;82:1882-91. (FC)
12. Zielinski CE, Mele F, Aschenbrenner D, Jarrossay D, Ronchi F, Gattorno M, et al. Pathogen-induced human T_H17 cells produce IFN- γ or IL-10 and are regulated by IL-1 β . *Nature.* 2012;484:514-8. (FC)
13. Sutton KM, Hu T, Wu Z, Siklodi B, Vervelde L, Kaiser P. The functions of the avian receptor activator of NF- κ B ligand (RANKL) and its receptors, RANK and osteoprotegerin, are evolutionarily conserved. *Dev Comp Immunol.* 2015;51:170-84. (FC)
14. Raats CJ, van den Born J, Bakker MA, Oppers-Walgreen B, Pisa BJ, Dijkman HB, et al. Expression of agrin, dystroglycan, and utrophin in normal renal tissue and in experimental glomerulopathies. *Am J Pathol.* 2000;156:1749-65. (IHC-FS, EM)
15. Poosti F, Bansal R, Yazdani S, Prakash J, Post E, Klok P, et al. Selective delivery of IFN- γ to renal interstitial myofibroblasts: a novel strategy for the treatment of renal fibrosis. *FASEB J.* 2015;29:1029-42. (IHC-FS)
16. Huizinga R, van der Star BJ, Kipp M, Jong R, Gerritsen W, Clamer T, et al. Phagocytosis of neuronal debris by microglia is associated with neuronal damage in multiple sclerosis. *Glia.* 2012;60:422-31. (IHC-PS)
17. Kreijtz JH, Wiersma LC, De Gruyter HL, Vogelzang-van Trierum SE, van Amerongen G, Stittelaar KJ, et al. A single immunization with modified vaccinia virus Ankara-based influenza virus H7 vaccine affords protection in the influenza A(H7N9) pneumonia ferret model. *J Infect Dis.* 2015;211:791-800. (IHC-PS)
18. Albrecht B, Woisetschlager M, Robertson MW. Export of the high affinity IgE receptor from the endoplasmic reticulum depends on a glycosylation-mediated quality control mechanism. *J Immunol.* 2000;165:5686-94. (ICC)
19. Haggqvist B, Hultman P. Effects of deviating the Th2-response in murine mercury-induced autoimmunity towards a Th1-response. *Clin Exp Immunol.* 2003;134:202-9. (ICC)
20. Mia MM, Bank RA. Paracrine factors of human amniotic fluid-derived mesenchymal stem cells show strong anti-fibrotic properties by inhibiting myofibroblast differentiation and collagen synthesis. *J Stem Cell Res Ther.* 2015;5:5. (ICC)
21. Santoro L, Reboul A, Kerblat I, Drouot C, Colomb MG. Monoclonal IgG as antigens: reduction is an early intracellular event of their processing by antigen-presenting cells. *Int Immunol.* 1996;8:211-9. (WB)
22. Attramadal CG, Halstensen TS, Dhakal HP, Ulekleiv CH, Boysen ME, Nesland JM, et al. High nuclear SOX2 expression is associated with radiotherapy response in small (T1/T2) oral squamous cell carcinoma. *J Oral Pathol Med.* 2015;44:515-22. (WB)
23. Konitsiotis AD, Jovanovic B, Ciepla P, Spitaler M, Lanyon-Hogg T, Tate EW, et al. Topological analysis of Hedgehog acyltransferase, a multipalmitoylated transmembrane protein. *J Biol Chem.* 2015;290:3293-307. (WB)
24. Sun Y, Peng I, Senger K, Hamidzadeh K, Reichelt M, Baca M, et al. Critical role of activation induced cytidine deaminase in experimental autoimmune encephalomyelitis. *Autoimmunity.* 2013;46:157-67. (Surface Plasmon Resonance)
25. Chmielewski M, Hombach AA, Abken H. CD28 cosignalling does not affect the activation threshold in a chimeric antigen receptor-redirectioned T-cell attack. *Gene Ther.* 2011;18:62-72. (Purification)

Texas Red® is a registered trademark of Molecular Probes, Inc.

Spectral Red® is a registered trademark of Southern Biotechnology Associates, Inc.

Spectral Red® is a PE/CY5 tandem conjugate.

Cy® is a registered trademark of GE Healthcare.

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.