




Rat Anti-Mouse IgE

Cat. No.	Format	Size
1130-01	Purified (UNLB)	0.5 mg
1130-02	Fluorescein (FITC)	0.5 mg
1130-04	Alkaline Phosphatase (AP)	1.0 mL
1130-05	Horseradish Peroxidase (HRP)	1.0 mL
1130-08	Biotin (BIOT)	0.5 mg
1130-09	R-phycoerythrin (PE)	0.1 mg
1130-09L	R-phycoerythrin (PE)	0.2 mg
1130-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg



Paraffin embedded mouse small intestine section was stained with Rat Anti-Mouse IgE-UNLB (SB Cat. No. 1130-01; right) followed by an HRP conjugated secondary antibody, DAB, and hematoxylin.

Overview

Clone	23G3
Isotype	Rat IgG ₁ κ
Immunogen	Monoclonal anti-DNP mouse IgE
Specificity	Mouse IgE

Applications

ELISA – Quality tested ¹⁻¹¹
 FLISA – Quality tested
 FC – Reported in literature ¹²⁻¹⁶
 IHC-FS – Reported in literature ^{17,18}
 IHC-PS ²³
 IHC-WM – Reported in literature ¹⁹
 WB – Reported in literature ^{5,21}
 ELISPOT – Reported in literature ^{1,20}
 Neut – Reported in literature ²²

Working Dilutions

ELISA	AP conjugate	1:500 – 1:1,000
	HRP conjugate	1:1,000 – 1:2,000
	BIOT conjugate	1:5,000 – 1:10,000
FLISA	FITC conjugate	1:200 – 1:400
	PE conjugate	≤ 1 µg/mL

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL or 0.2 mg in 2.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Petit-Frere C, Dugas B, Braquet P, Mencia-Huerta JM. Interleukin-9 potentiates the interleukin-4-induced IgE and IgG1 release from murine B lymphocytes. *Immunology*. 1993;79:146-51. (ELISA, ELISPOT)
2. Takahashi I, Marinaro M, Kiyono H, Jackson RJ, Nakagawa I, Fujihashi K, et al. Mechanisms for mucosal immunogenicity and adjuvancy of *Escherichia coli* labile enterotoxin. *J Infect Dis*. 1996;173:627-35. (ELISA)
3. Kasprovicz DJ, Kohm AP, Berton MT, Chruscinski AJ, Sharpe A, Sanders VM, et al. Stimulation of the B cell receptor, CD86 (B7-2), and the β_2 -adrenergic receptor intrinsically modulates the level of IgG1 and IgE produced per B cell. *J Immunol*. 2000;165:680-90. (ELISA)
4. Häggqvist B, Hultman P. Effects of deviating the Th2-response in murine mercury-induced autoimmunity towards a Th1-response. *Clin Exp Immunol*. 2003;134:202-9. (ELISA)
5. Lehrer SB, Reish R, Fernandes J, Gaudry P, Dai G, Reese G. Enhancement of murine IgE antibody detection by IgG removal. *J Immunol Methods*. 2004;284:1-6. (ELISA, WB)
6. Ramalingam TR, Pesce JT, Sheikh F, Cheever AW, Mentink-Kane MM, Wilson MS, et al. Unique functions of the type II interleukin 4 receptor identified in mice lacking the interleukin 13 receptor $\alpha 1$ chain. *Nat Immunol*. 2008;9:25-33. (ELISA)
7. Li F, Wang L, Jin X, Yan C, Jiang S, Shen X. The immunologic effect of TGF- β 1 chitosan nanoparticle plasmids on ovalbumin-induced allergic BALB/c mice. *Immunobiology*. 2009;214:87-99. (ELISA)
8. Kamphuis T, Meijerhof T, Stegmann T, Lederhofer J, Wilschut J, de Haan A. Immunogenicity and protective capacity of a virosomal respiratory syncytial virus vaccine adjuvanted with monophosphoryl lipid A in mice. *PLoS One*. 2012;7(5):e36812. (ELISA)
9. Futatsugi-Yumikura S, Matsushita K, Fukuoka A, Takahashi S, Yamamoto N, Yonehara S, et al. Pathogenic T_H2-type follicular helper T cells contribute to the development of lupus in Fas-deficient mice. *Int Immunol*. 2014;26:221-31. (ELISA)
10. Stefka AT, Feehley T, Tripathi P, Qiu J, McCoy K, Mazmanian SK, et al. Commensal bacteria protect against food allergen sensitization. *Proc Natl Acad Sci USA*. 2014;111:13145-50. (ELISA)
11. Pucella JN, Yen W, Kim MV, van der Veecken J, Socci ND, Naito Y, et al. miR-182 is largely dispensable for adaptive immunity: lack of correlation between expression and function. *J Immunol*. 2015;194:2635-42. (ELISA)
12. Kuzin II, Snyder JE, Ugine GD, Wu D, Lee S, Bushnell T Jr, et al. Tetracyclines inhibit activated B cell function. *Int Immunol*. 2001;12:921-31. (FC)
13. Huber M, Kalis C, Keck S, Jiang Z, Georgel P, Du X, et al. R-form LPS, the master key to the activation of TLR4/MD-2-positive cells. *Eur J Immunol*. 2006;36:701-11. (FC)
14. Seidl A, Panzer M, Voehringer D. Protective immunity against the gastrointestinal nematode *Nippostrongylus brasiliensis* requires a broad T-cell receptor repertoire. *Immunology*. 2011;134:214-23. (FC)
15. Kamijo S, Takeda H, Tokura T, Suzuki M, Inui K, Hara M, et al. IL-33-mediated innate response and adaptive immune cells contribute to maximum responses of protease allergen-induced allergic airway inflammation. *J Immunol*. 2013;190:4489-99. (FC)
16. Muto T, Fukuoka A, Kabashima K, Ziegler SF, Nakanishi K, Matsushita K, et al. The role of basophils and proallergic cytokines, TSLP and IL-33, in cutaneously sensitized food allergy. *Int Immunol*. 2014;26:539-49. (FC)
17. Brown JK, Knight PA, Pemberton AD, Wright SH, Pate JA, Thornton EM, et al. Expression of integrin- α_E by mucosal mast cells in the intestinal epithelium and its absence in nematode-infected mice lacking the transforming growth factor- β_1 -activating integrin $\alpha_V\beta_6$. *Am J Pathol*. 2004;165:95-106. (IHC-FS)
18. Inman CF, Rees LE, Barker E, Haverson K, Stokes CR, Bailey M. Validation of computer-assisted, pixel-based analysis of multiple-colour immunofluorescence histology. *J Immunol Methods*. 2005;302:156-67. (IHC-FS)
19. Pochanke V, Koller S, Dayer R, Hatak S, Ludewig B, Zinkernagel RM, et al. Identification and characterization of a novel antigen from the nematode *Nippostrongylus brasiliensis* recognized by specific IgE. *Eur J Immunol*. 2007;37:1275-84. (IHC-WM)
20. Padro CJ, Shawler TM, Gormley MG, Sanders VM. Adrenergic regulation of IgE involves modulation of CD23 and ADAM10 expression on exosomes. *J Immunol*. 2013;191:5383-97. (ELISPOT)
21. Ambrosi G, Ghezzi C, Sepe S, Milanese C, Payan-Gomez C, Bombardieri CR, et al. Bioenergetic and proteolytic defects in fibroblasts from patients with sporadic Parkinson's disease. *Biochim Biophys Acta*. 2014;1842:1385-94. (WB)
22. Dakhama A, Lee Y, Ohnishi H, Jing X, Balhorn A, Takeda K, et al. Virus-specific IgE enhances airway responsiveness on reinfection with respiratory syncytial virus in newborn mice. *J Allergy Clin Immunol*. 2009;123:138-45. (Neut)
23. SouthernBiotech published data (IHC-PS)