

# **Mycoplasma Detection Kit**

## **User Guide**

**Cat. No. 13100-01**

**50 Reactions**

*For Research Use Only*

## Description

---

The SouthernBiotech Mycoplasma Detection Kit is designed to specifically detect potential *Mycoplasma* contamination in cell cultures. The kit incorporates polymerase chain reaction (PCR) to amplify the conserved 16S ribosomal RNA coding region within the *Mycoplasma* genome, thereby providing an extensive, highly sensitive, and efficient detection method. Carefully determined primer sequences cover three genera of Mycoplasmatales (*Mycoplasma*, *Acholeplasma*, and *Ureaplasma*) which allows the kit to detect over 95% of potential cell culture infections.

The PCR technology in the Mycoplasma Detection Kit is fast (results are typically obtained in less than 3 hours) and easy to use. The kit is also highly sensitive and can detect as little as 2-5 femtograms of *Mycoplasma* DNA in 100  $\mu$ L of test sample supernatant. Eukaryotic and bacterial DNA from cell culture supernatant is not amplified by the kit.

A sample cell line infected with *Mycoplasma* will generate a PCR product between ~448 bp to ~611 bp on an agarose gel, depending on the type of *Mycoplasma* present. A positive control (*M. orale*, 503 bp) is included to validate that the PCR amplification process has occurred as well as to confirm the size of the PCR product obtained in test samples. An internal control is also provided to eliminate potential false negatives associated with PCR inhibitors.

## Kit Components

---

- User guide
- Primer set and nucleotides (lyophilized) - Blue Cap
  - Primer/dNTP mix (deoxynucleotide triphosphates including dATP, dCTP, dGTP, and dUTP)
- Sterile PCR 10X reaction buffer (500  $\mu$ L) - White Cap
  - 100 mM Tris-HCl, pH 8.5
  - 750 mM KCl
  - 30 mM MgCl<sub>2</sub>
- Positive Control DNA (lyophilized) - Yellow Cap
  - Non-infectious PCR product from *M. orale*
  - Yields a 503 bp band
- Internal Control DNA (lyophilized) - Green Cap
  - Non-infectious plasmid DNA including *Mycoplasma*-specific primer sequences
  - Yields a 270 bp band

## Required Materials

---

- Thermal cycler
- Taq DNA polymerase
- DNA ladder
- PCR reaction tubes
- DNA-free water

## Strains Amplified

---

Strain	Base Pairs (bp)	Strain	Base Pairs (bp)
<i>A. laidlawii</i>	611	<i>M. hyorhinis</i>	525
<i>M. arginini</i>	448	<i>M. hyosynoviae</i>	471
<i>M. arthritidis</i>	486	<i>M. opalescens</i>	573
<i>M. bovis</i>	605	<i>M. orale</i>	503
<i>M. cloacale</i>	459	<i>M. pirum</i>	533
<i>M. falconis</i>	458	<i>M. pneumonia</i>	493
<i>M. faucium</i>	479	<i>M. salivarium</i>	482
<i>M. fermentans</i>	579	<i>M. synoviae</i>	572
<i>M. genitalium</i>	464	<i>U. urealyticum</i>	558
<i>M. hominis</i>	448		

## Handling and Storage

---

The kit is stable for the period shown on the label when stored as directed. Upon reconstitution of the primer/dNTP mix, the positive control, and the internal control, store below -20°C. Avoid freeze / thaw cycles.

## Suggested Protocol

- Reagent Preparation
  - Centrifuge all kit component tubes to ensure that material has completely settled to the bottom
  - Reconstitute primer/dNTP mix, positive, and internal control using DNA-free water as follows -
    - Primer/dNTP mix - 260  $\mu$ L
    - Positive control - 200  $\mu$ L
    - Internal control - 200  $\mu$ L
  - Vortex for 5 seconds to ensure thorough mixing and equilibrate at room temperature for 5 minutes
  - Vortex, then centrifuge again
  - Reconstituted reagents should be stored below  $-20^{\circ}\text{C}$
- Sample Preparation
  - Transfer 100  $\mu$ L of cell culture supernatant\* to a sterile 200  $\mu$ L PCR tube; ensure lid has been tightly sealed to prevent evaporation
  - Heat the sample supernatant at  $95^{\circ}\text{C}$  for 5 minutes
  - Quickly spin the sample supernatant at maximum speed for 5 seconds to remove cell debris; the supernatant is now ready to add to the PCR master mix

\*This kit is specifically designed for *Mycoplasma* detection from cell culture supernatant. For serum or cell lysate samples, DNA extraction is recommended prior to testing to avoid PCR inhibition.

- Thermal Cycling Program
  - Incubation times are dependent on the type of polymerase used; the template described below has been optimized for hot start Taq polymerase
    - 1 cycle             $95^{\circ}\text{C}$  for 5 minutes
    - 35 cycles         $95^{\circ}\text{C}$  for 30 seconds  
                       $58^{\circ}\text{C}$  for 40 seconds  
                       $72^{\circ}\text{C}$  for 1 minute
    - Cool down to  $4^{\circ}\text{C}$

- PCR Master Mix
  - Total volume per reaction is 50  $\mu$ L
    - When preparing reactions, calculations should also include positive and negative controls
    - Add 48  $\mu$ L of master mix to each tube
    - Contents of master mix are provided below -

Component	1 Reaction ( $\mu$ L)	10 Reactions ( $\mu$ L)	50 Reactions ( $\mu$ L)
Water	35.6	356	1780
10X reaction buffer	5	50	250
Primer/dNTP mix	5	50	250
Internal control	2	20	100
Taq polymerase (5U/ $\mu$ L)*	0.4	4	20

Note - If using a polymerase other than Taq, use the PCR buffer supplied with the enzyme and the final  $\text{MgCl}_2$  concentration must be adjusted to 3.0 mM


- Add 2  $\mu$ L of DNA-free water as a negative control into the appropriate PCR reaction tube
- Add 2  $\mu$ L of reconstituted positive control DNA supplied in the kit into the appropriate PCR reaction tube
- Add 2  $\mu$ L of heat-treated test sample into the appropriate PCR reaction tube
- Run according to the thermal cycling program
- Agarose Gel Electrophoresis
  - Use a 1% - 1.5% agarose gel containing 0.5  $\mu\text{g}/\text{mL}$  ethidium bromide
  - Load 15  $\mu$ L - 20  $\mu$ L of each PCR reaction mixed with bromophenol blue loading buffer into each well
  - Run at 100 V for 20 min
  - Detect PCR product bands using UV box

Note - Ethidium bromide is a known mutagen and should be handled as a hazardous chemical; wear gloves when handling

- Gel Evaluation

- Samples containing *Mycoplasma* infection will contain a band (or multiple bands) between ~448 bp to ~611 bp
- Samples containing the supplied internal control DNA will contain a distinct 270 bp PCR product that is clearly distinguishable from the ~448 bp to ~611 bp band(s) found in positive samples (see sample data, lane 7)
  - The internal control indicates a successfully performed PCR reaction
  - PCR inhibition may have occurred if the internal control band disappears in some samples, but the band is present in the PCR reaction of the negative control (water added instead of test sample)
    - If the PCR of a sample is inhibited, the inhibitors can be easily removed by performing a DNA extraction
- If the cell culture is heavily contaminated with *Mycoplasma*, amplification of the ~448 bp to ~611 bp product(s) may diminish or completely eliminate the 270-bp internal control product (see sample data, lane 6)
- Low intensity bands smaller than 100 bp indicate the presence of non-specific, self-annealing primers; this does not indicate a positive result and will not affect the precision of the test

## Sample Data



Lane 1	<i>E. coli</i> Genomic DNA	Lane 5	Inhibited Sample
Lane 2	Hybridoma Cell DNA	Lane 6	Strongly Contaminated Sample
Lane 3	Negative Control (H <sub>2</sub> O)	Lane 7	Weakly Contaminated Sample
Lane 4	Positive Control	Lane 8	100 bp DNA Ladder

## References

- Lamb SA, Rahman MM, McFadden G. Recombinant myxoma virus lacking all poxvirus ankyrin-repeat proteins stimulates multiple cellular anti-viral pathways and exhibits a severe decrease in virulence. *Virology*. 2014;464-5:134-45.
- Frampton GM, Ali SM, Rosenzweig M, Chmielecki J, Lu X, Bauer TM, et al. Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov*. 2015;5:850-9.
- Modur V, Joshi P, Nie D, Robbins KT, Khan AU, Rao K. CD24 expression may play a role as a predictive indicator and a modulator of cisplatin treatment response in head and neck squamous cellular carcinoma. *PLoS One*. 2016;11(6):e0156651.
- Campos-Gomez J, Ahmad F, Rodriguez E, Saeed MF. A novel cell-based assay to measure activity of Venezuelan equine encephalitis virus nsP2 protease. *Virology*. 2016;496:77-89.
- Wang X, Shaw DK, Hammond HL, Sutterwala FS, Rayamajhi M, Shirey KA, et al. The prostaglandin E<sub>2</sub>-EP3 receptor axis regulates Anaplasma phagocytophilum-mediated NLR4 inflammasome activation. *PLoS Pathog*. 2016;12(8):e01005803.
- Baird JR, Feng Z, Xiao HD, Friedman D, Cottam B, Fox BA, et al. STING expression and response to treatment with STING ligands in premalignant and malignant disease. *PLoS One*. 2017;12(11):e0187532.
- Melvin JA, Gaston JR, Phillips SN, Springer MJ, Marshall CW, Shanks RM, et al. Pseudomonas aeruginosa contact-dependent growth inhibition plays dual role in host-pathogen interactions. *mSphere*. 2017;2:e00336-17.
- Wagner RD, Johnson SJ, Danielsen ZY, Lim J, Mudalige T, Linder S. Polyethylene glycol-functionalized poly (lactic acid-co-glycolic acid) and graphene oxide nanoparticles induce pro-inflammatory and apoptotic responses in Candida albicans-infected vaginal epithelial cells. *PLoS One*. 2017;12(4):e0175250.
- McGee D, Smith A, Poncil S, Patterson A, Bernstein AI, Racicot K. Cervical HSV-2 infection causes cervical remodeling and increases risk for ascending infection and preterm birth. *PLoS One*. 2017;12(11):e0188645.
- Sprague L, Lee JM, Hutzen BJ, Wang P, Chen C, Conner J, et al. High mobility group box 1 influences HSV1716 spread and acts as an adjuvant to chemotherapy. *Viruses*. 2018;10:132.
- Crittenden MR, Zebertavage L, Kramer G, Bambina S, Friedman D, Troesch V, et al. Tumor cure by radiation therapy and checkpoint inhibitors depends on pre-existing immunity. *Sci Rep*. 2018;8:7012.
- Razidlo GL, Burton KM, McNiven MA. Interleukin-6 promotes pancreatic cancer cell migration by rapidly activating the small GTPase CDC42. *J Biol Chem*. 2018;293:11143-53.
- Pishas KI, Drenberg CD, Taslim C, Theisen ER, Johnson KM, Saund RS, et al. Therapeutic targeting of KDM1A/LSD1 in Ewing sarcoma with SP-2509 engages the endoplasmic reticulum stress response. *Mol Cancer Ther*. 2018;17:1902-16.
- Bryant JD, Sweeney SR, Sentandreu E, Shin M, Ipas H, Xhemalce B, et al. Deletion of the neural tube defect-associated gene Mthfd11 disrupts one-carbon and central energy metabolism in mouse embryos. *J Biol Chem*. 2018;293:5821-33.