




## Hamster Anti-Mouse CD3 $\epsilon$

Cat. No.	Format	Size
1531-01	Purified (UNLB)	0.5 mg
1531-02	Fluorescein (FITC)	0.5 mg
1531-08	Biotin (BIOT)	0.5 mg
1531-09	R-phycoerythrin (PE)	0.1 mg
1531-11	Allophycocyanin (APC)	0.1 mg
1531-13	Spectral Red <sup>®</sup> (SPRD)	0.1 mg
1531-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg
1531-15	Cyanine 5 (CY5)	0.1 mg
1531-18	Allophycocyanin-Cyanine 5.5 (APC/CY5.5)	0.1 mg



BALB/c mouse splenocytes were stained with Hamster Anti-Mouse CD3 $\epsilon$ -APC (SB Cat. No. 1531-11) and Rat Anti-Mouse CD19-FITC (SB Cat. No. 1575-02).

### Overview

<b>Clone</b>	500-A2
<b>Isotype</b>	Hamster (Syrian) IgG <sub>2</sub>
<b>Immunogen</b>	Immune complexes isolated from C6VL-BS thymoma cells
<b>Specificity</b>	Mouse CD3 $\epsilon$ ; Mr 25 kDa
<b>Alternate Name(s)</b>	CD3 epsilon

### Description

CD3 $\epsilon$ , a member of the immunoglobulin superfamily of cell surface receptors, is comprised of five invariable chains ranging in size from 16-28 kDa and is closely associated with the T cell antigen receptor (TCR). CD3 $\epsilon$  is expressed on all T cells of all mouse strains. CD3 plays a major role in signaling during antigen recognition, leading to T-cell activation. 500-A2 recognizes an epitope on the 25 kDa  $\epsilon$  chain of the CD3/TCR complex of mouse strains tested.

### Applications

FC – Quality tested <sup>1</sup>  
 IHC-FS – Reported in literature <sup>3</sup>  
 IP – Reported in literature <sup>1</sup>  
 Activ – Reported in literature <sup>1,2,4</sup>

### Working Dilutions

<b>Flow Cytometry</b>	FITC conjugate	$\leq 3 \mu\text{g}/10^6$ cells
	BIOT conjugate	$\leq 1 \mu\text{g}/10^6$ cells
	PE, APC, and APC/CY5.5 conjugates	$\leq 0.3 \mu\text{g}/10^6$ cells
	SPRD and CY5 conjugates	$\leq 0.1 \mu\text{g}/10^6$ cells
For flow cytometry, the suggested use of these reagents is in a final volume of 100 $\mu\text{l}$		

**Other Applications** Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: [info@southernbiotech.com](mailto:info@southernbiotech.com) • Website: [www.southernbiotech.com](http://www.southernbiotech.com)

## Handling and Storage

---

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) and allophycocyanin (APC) conjugates are supplied as 0.1 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral Red<sup>®</sup> (SPRD) and allophycocyanin-Cyanine 5.5 (APC/CY5.5) conjugates are supplied as 0.1 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- The Cyanine 5 (CY5) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

---

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

## References

---

1. Allison JP, Havran WL, Poenie M, Kimura J, DeGraffenreid L, Ajami M, et al. Expression and function of CD3 on murine thymocytes. In: Davis MM, Kappler J, editors. *The T-Cell Receptor, UCLA Symposia*, 73<sup>rd</sup> Edition. New York: Alan R. Liss; 1988. p.33-45. (Immunogen, FC, IP, Activ)
2. Swallow MM, Wallin JJ, Sha WC. B7h, a novel costimulatory homolog of B7.1 and B7.2, is induced by TNF $\alpha$ . *Immunity*. 1999;11:423-32. (Activ)
3. Hörnquist CE, Lu X, Rogers-Fani PM, Rudolph U, Shappell S, Birnbaumer L, et al. *Gai2*-deficient mice with colitis exhibit a local increase in memory CD4<sup>+</sup> T cells and proinflammatory Th1-type cytokines. *J Immunol*. 1997;158:1068-77. (IHC-FS)
4. Portoles P, Rojo J, Golby A, Bonneville M, Gromkowski S, Greenbaum L, et al. Monoclonal antibodies to murine CD3 $\epsilon$  define distinct epitopes one of which may interact with CD4 during T cell activation. *J Immunol*. 1989;142:4169-75. (Activ)

Spectral Red<sup>®</sup> is a registered trademark of Southern Biotechnology Associates, Inc.

Spectral Red<sup>®</sup> is a PE/CY5 tandem conjugate.

Cy<sup>®</sup> is a registered trademark of GE Healthcare.

Alexa Fluor<sup>®</sup> 488, 647, 700 and Pacific Blue<sup>™</sup> are provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Southern Biotechnology Associates, Inc., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.