




Rat Anti-Mouse CD11b

Cat. No.	Format	Size
1560-01	Purified (UNLB)	0.5 mg
1560-02S	Fluorescein (FITC)	0.1 mg
1560-02	Fluorescein (FITC)	0.5 mg
1560-08	Biotin (BIOT)	0.5 mg
1560-09	R-phycoerythrin (PE)	0.1 mg
1560-09L	R-phycoerythrin (PE)	0.2 mg
1560-11	Allophycocyanin (APC)	0.1 mg
1560-13	Spectral Red [®] (SPRD)	0.1 mg
1560-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg
1560-15	Cyanine 5 (CY5)	0.1 mg
1560-17	R-phycoerythrin-Cyanine 7 (PE/CY7)	0.1 mg
1560-19	Allophycocyanin-Cyanine 7 (APC/CY7)	0.1 mg
1560-27	Alexa Fluor [®] 700 (AF700)	0.1 mg
1560-30	Alexa Fluor [®] 488 (AF488)	0.1 mg



BALB/c mouse bone marrow cells were stained with Rat Anti-Mouse CD11b-APC (SB Cat. No. 1560-11).

Overview

Clone	3A33
Isotype	Rat (Lewis) IgG _{2a} κ
Immunogen	Peritoneal macrophages from B6D2 hybrid mice
Specificity	Mouse CD11b; Mr 170 kDa
Alternate Name(s)	Integrin α _M , Mac-1α, Mac-1, Mo1, CR3, C3biR

Description

CD11b (integrin α_M subunit) combines with CD18 (integrin β₂ subunit) to form the integrin Mac-1, also known as complement receptor 3 (CR3), which mediates adhesion to C3bi and ICAM-1 (CD54). Mac-1 is expressed at varying levels on granulocytes, macrophages, dendritic cells, NK cells, and B-1 cells in the peritoneal and pleural cavities. In addition to its role in binding C3bi on opsonized targets and mediation of the subsequent ingestion process, Mac-1 is important as an adhesion molecule in the transendothelial migration of monocytes and neutrophils.

Applications

FC – Quality tested^{1,4-13}
 IHC-FS – Reported in literature¹⁴
 ICC – Reported in literature²
 IP – Reported in literature¹
 Block – Reported in literature³

Working Dilutions

Flow Cytometry	Purified (UNLB) antibody	≤ 1 μg/10 ⁶ cells
	FITC, BIOT, and AF488 conjugates	≤ 1 μg/10 ⁶ cells
	PE, APC, SPRD, CY5, PE/CY7, APC/CY7, and AF700 conjugates	≤ 0.3 μg/10 ⁶ cells

For flow cytometry, the suggested use of these reagents is in a final volume of 100 μL

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: info@southernbiotech.com • Website: www.southernbiotech.com

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg or 0.1 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL or 0.2 mg in 2.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The allophycocyanin (APC) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral Red[®] (SPRD), R-phycoerythrin-Cyanine 7 (PE/CY7) and allophycocyanin-Cyanine 7 (APC/CY7) conjugates are supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- The Cyanine 5 (CY5) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The Alexa Fluor[®] 488 (AF488) and Alexa Fluor[®] 700 (AF700) conjugates are supplied as 0.1 mg in 0.2 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Martin A, Le Corre R, Pellen P, Bourel D, Merdrignac G, Genetet B, et al. A monoclonal antibody (3A33) that reacts with a mouse-specific epitope of Mac-1 antigen. *Tissue Antigens*. 1986;28:15-23. (Immunogen, IP, FC)
2. Wang M, Shakhathreh MK, James D, Liang S, Nishiyama S, Yoshimura F, et al. Fimbrial proteins of porphyromonas gingivalis mediate in vivo virulence and exploit TLR2 and complement receptor 3 to persist in macrophages. *J Immunol*. 2007;179:2349-58. (ICC)
3. Collet B, Martin A, Merdrignac G, Genetet B, Toujas L. Modulation of expression of mouse macrophage surface antigens by monoclonal antibodies. *Immunology*. 1985;56:543-8. (Block)
4. Jang Y, Collector MI, Baylin SB, Diehl AM, Sharkis SJ. Hematopoietic stem cells convert into liver cells within days without fusion. *Nat Cell Biol*. 2004;6:532-9. (FC)
5. Lee Y, Kim J, Choi I, Kim H, Cho S, et al. Enhanced antitumor effect of oncolytic adenovirus expressing interleukin-12 and B7-1 in an immunocompetent murine model. *Clin Cancer Res*. 2006;12:5859-68. (FC)
6. Choi K, Kim J, Lee Y, Kim J, Suh B, Kim H, et al. Concurrent delivery of GM-CSF and B7-1 using an oncolytic adenovirus elicits potent antitumor effect. *Gene Ther*. 2006;13:1010-20. (FC)
7. Pevsner-Fischer M, Morad V, Cohen-Sfady M, Rousso-Noori L, Zanin-Zhorov A, Cohen S, et al. Toll-like receptors and their ligands control mesenchymal stem cell functions. *Blood*. 2007;109:1422-32. (FC)
8. Morad V, Pevsner-Fischer M, Barnees S, Samokovlisky A, Rousso-Noori L, Rosenfeld R, et al. The myelopoietic supportive capacity of mesenchymal stromal cells is uncoupled from multipotency and is influenced by lineage determination and interference with glycosylation. *Stem Cells*. 2008;26:2275-86. (FC)
9. Avagyan S, Glouchkova L, Choi J, Snoeck H. A quantitative trait locus on chromosome 4 affects cycling of hematopoietic stem and progenitor cells through regulation of TGF-β2 responsiveness. *J Immunol*. 2008;181:5904-11. (FC)
10. You Y, Zhao H, Wang Y, Carter RH. Cutting edge: Primary and secondary effects of CD19 deficiency on cells of the marginal zone. *J Immunol*. 2009;182:7343-47. (FC)
11. Shainer R, Azar Y, Almogi-Hazan O, Bringer R, Compton SR, Paidas MJ, et al. Immune regulation and oxidative stress reduction by preimplantation factor following syngeneic or allogeneic bone marrow transplantation. *Conference Papers in Medicine*. 2013;11:1-8. (FC)
12. Azar Y, Shainer R, Almogi-Hazan O, Bringer R, Compton SR, Paidas MJ, et al. Preimplantation factor reduces graft-versus-host disease by regulating immune response and lowering oxidative stress (murine model). *Biol Blood Marrow Transplant*. 2013;19:519-28. (FC)
13. Knight DA, Ngiew SF, Li M, Parmenter T, Mok S, Cass A, et al. Host immunity contributes to the anti-melanoma activity of BRAF inhibitors. *J Clin Invest*. 2013;123:1371-81. (FC)
14. Everett J, Turner K, Cai Q, Gordon V, Whiteley M, Rumbaugh K. Arginine is a critical substrate for the pathogenesis of *Pseudomonas aeruginosa* in burn wound infections. *MBio*. 2017;8:e02160-16. (IHC-FS)

Spectral Red[®] is a registered trademark of Southern Biotechnology Associates, Inc.

Spectral Red[®] is a PE/CY5 tandem conjugate.

Cy[®] is a registered trademark of GE Healthcare.

Alexa Fluor[®] 488, 647, 700 and Pacific Blue[™] are provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Southern Biotechnology Associates, Inc., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.