




## Hamster Anti-Mouse CD103

Cat. No.	Format	Size
1810-01	Purified (UNLB)	0.5 mg
1810-02	Fluorescein (FITC)	0.5 mg
1810-08	Biotin (BIOT)	0.5 mg
1810-09	R-phycoerythrin (PE)	0.1 mg
1810-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg
1810-26	Pacific Blue™ (PACBLU)	0.1 mg
1810-30	Alexa Fluor® 488 (AF488)	0.1 mg
1810-31	Alexa Fluor® 647 (AF647)	0.1 mg



CD-1 mouse mesenteric lymph node cells were stained with Hamster Anti-Mouse CD103-FITC (SB Cat. No. 1810-02) and Rat Anti-Mouse CD3ε-PE (SB Cat. No. 1535-09).

### Overview

<b>Clone</b>	2E7
<b>Isotype</b>	Hamster (Armenian) IgG <sub>2</sub>
<b>Immunogen</b>	C57BL/6J mouse intestinal intraepithelial lymphocytes (IEL)
<b>Specificity</b>	Mouse CD103; Mr 150 & 25 kDa
<b>Alternate Name(s)</b>	Integrin $\alpha_{IEL}$ , $\alpha_E$ integrin, ITGAE

### Description

CD103 is a member of the integrin series of adhesion molecules. This antigen defines a developmentally important subset of T cells, namely mucosal T cells including all IEL (intraepithelial lymphocytes) and ~20% of lamina propria T cells. Expression of CD103 is more restricted outside these mucosal organs appearing at lower levels on T cell subsets of the lymph node, dendritic epidermis, and periphery. In non-epithelial CD103<sup>+</sup> T cells there is a bias toward expression on CD8<sup>+</sup> cells. The 2E7 monoclonal antibody exhibits antigen immunoprecipitation patterns similar if not identical to the rat anti-mouse CD103 monoclonal antibody M290. 2E7 is reported to have signal-inducing activity in a redirected lysis assay and to costimulate IEL and CD8<sup>+</sup> lymph node cells in conjunction with anti-TCR *in vitro* which runs counter to the usual inhibition of CTL by anti-integrins. This costimulatory activity is also shared by the monoclonal antibody M290. The property of costimulation is not unique to the mouse since the combination of anti-human  $\alpha_{IEL}$  plus anti-TCR will produce similar proliferation in human cells with the exception that CD8<sup>+</sup> cells are not stimulated.

### Applications

FC – Quality tested <sup>2</sup>  
 IP – Reported in literature <sup>2</sup>  
 IHC-FS – Reported in literature <sup>2</sup>  
 Activ – Reported in literature <sup>3</sup>

### Working Dilutions

<b>Flow Cytometry</b>	FITC, BIOT, PACBLU, and AF488 conjugates	≤ 1 μg/10 <sup>6</sup> cells
	PE and AF647 conjugates	≤ 0.2 μg/10 <sup>6</sup> cells
For flow cytometry, the suggested use of these reagents is in a final volume of 100 μL		

**Other Applications** Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: [info@southernbiotech.com](mailto:info@southernbiotech.com) • Website: [www.southernbiotech.com](http://www.southernbiotech.com)

## Handling and Storage

---

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- The Alexa Fluor<sup>®</sup> 488 (AF488), Alexa Fluor<sup>®</sup> 647 (AF647), and Pacific Blue<sup>™</sup> (PACBLU) conjugates are supplied as 0.1 mg in 0.2 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

---

Some reagents contain sodium azide. Please refer to product specific SDS.

## References

---

1. Goodman T, Lefrancois L. Intraepithelial lymphocytes. Anatomical site, not T cell receptor form, dictates phenotype and function. *J Exp Med.* 1989;170:1569-81. (Immunogen)
2. Lefrançois L, Barrett TA, Havran WL, Puddington L. Developmental expression of the  $\alpha_{IEL}\beta_7$  integrin on T cell receptor  $\gamma\delta$  and T cell receptor  $\alpha\beta$  T cells. *Eur J Immunol.* 1994;24:635-40. (IP, IHC-FS, FC)
3. Müller S, Jungo M, Aichele P, Mueller C. CD5<sup>-</sup> CD8 $\alpha\beta$  intestinal intraepithelial lymphocytes (IEL) are induced to express CD5 upon antigen-specific activation: CD5<sup>-</sup> and CD5<sup>+</sup> CD8 $\alpha\beta$  IEL do not represent separate T cell lineages. *Eur J Immunol.* 1997;27:1756-61. (Activ)

Alexa Fluor<sup>®</sup> 488, 647, 700 and Pacific Blue<sup>™</sup> are provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Southern Biotechnology Associates, Inc., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.