




Rat Anti-Mouse Ly-6G/Ly-6C

Cat. No.	Format	Size
1900-01	Purified (UNLB)	0.5 mg
1900-02	Fluorescein (FITC)	0.5 mg
1900-08	Biotin (BIOT)	0.5 mg
1900-09	R-phycoerythrin (PE)	0.1 mg
1900-09L	R-phycoerythrin (PE)	0.2 mg
1900-11	Allophycocyanin (APC)	0.1 mg
1900-13	Spectral Red® (SPRD)	0.1 mg
1900-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg
1900-15	Cyanine 5 (CY5)	0.1 mg
1900-16	R-phycoerythrin-Cyanine 5.5 (PE/CY5.5)	0.1 mg
1900-17	R-phycoerythrin-Cyanine 7 (PE/CY7)	0.1 mg
1900-19	Allophycocyanin-Cyanine 7 (APC/CY7)	0.1 mg
1900-26	Pacific Blue™ (PACBLU)	0.1 mg
1900-27	Alexa Fluor® 700 (AF700)	0.1 mg
1900-30	Alexa Fluor® 488 (AF488)	0.1 mg
1900-31	Alexa Fluor® 647 (AF647)	0.1 mg



BALB/c mouse bone marrow cells were stained with Rat Anti-Mouse Ly-6G/Ly-6C-PE/CY7 (SB Cat. No. 1900-17).

Overview

Clone	RB6-8C5
Isotype	Rat IgG _{2b} K
Immunogen	Unknown
Specificity	Mouse Ly-6G/Ly-6C; Mr 21-25 kDa
Alternate Name(s)	Gr-1

Description

Ly-6G/Ly-6C, formerly known as the myeloid differentiation antigen Gr-1, is a GPI-anchored protein. In the bone marrow, the level of antigen expression is directly correlated with granulocyte differentiation and maturation. It is also transiently expressed on monocytes in the bone marrow. In the periphery, Ly-6G/Ly-6C is expressed predominantly on neutrophils.

Applications

FC – Quality tested ^{1,11,12}
 IHC-FS – Reported in literature ⁷⁻¹⁰
 IP – Reported in literature ¹
 CMCD – Reported in literature ²
 Depletion – Reported in literature ³⁻⁶

Working Dilutions

Flow Cytometry	FITC, BIOT, AF488, and PACBLU conjugates	≤ 1 µg/10 ⁶ cells
	PE and AF647 conjugates	≤ 0.1 µg/10 ⁶ cells
	SPRD, APC, CY5, PE/CY5.5, PE/CY7, APC/CY7, and AF700 conjugates	≤ 0.03 µg/10 ⁶ cells

For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL.

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL or 0.2 mg of 2.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The allophycocyanin (APC) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral Red[®] (SPRD), R-phycoerythrin-Cyanine 5.5 (PE/CY5.5), R-phycoerythrin-Cyanine 7 (PE/CY7) and allophycocyanin-Cyanine 7 (APC/CY7) conjugates are supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C immediately upon receipt.**
- The Cyanine 5 (CY5) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The Alexa Fluor[®] 488 (AF488), Alexa Fluor[®] 647 (AF647), Alexa Fluor[®] 700 (AF700), and Pacific Blue[™] (PACBLU) conjugates are supplied as 0.1 mg in 0.2 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Fleming TJ, Fleming ML, Malek TR. Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. *J Immunol.* 1993;151:2399-408. (FC, IP)
2. Brummer E, Sugar AM, Stevens DA. Immunological activation of polymorphonuclear neutrophils for fungal killing: studies with murine cells and blastomyces dermatitidis in vitro. *J Leukoc Biol.* 1984;36:505-20. (CMCD)
3. Franco JL, Ghosh P, Wiltout RH, Carter CR, Zea AH, Momozaki N, et al. Partial degradation of T-cell signal transduction molecules by contaminating granulocytes during protein extraction of splenic T cells from tumor-bearing mice. *Cancer Res.* 1995;55:3840-6. (Depletion)
4. Conlan JW, North RJ. Neutrophils are essential for early anti-Listeria defense in the liver, but not in the spleen or peritoneal cavity, as revealed by a granulocyte-depleting monoclonal antibody. *J Exp Med.* 1994;179:259-68. (Depletion)
5. Tumpey TM, Chen SH, Oakes JE. Neutrophil-mediated suppression of virus replication after herpes simplex virus type 1 infection of the murine cornea. *J Virol.* 1996;70:898-904. (Depletion)
6. Gong Y, Koh D. Neutrophils promote inflammatory angiogenesis via release of preformed VEGF in an in vivo corneal model. *Cell Tissue Res.* 2010;339:437-48. (Depletion)
7. Okuma T, Terasaki Y, Kaikita K, Kobayashi H, Kuziel WA, Kawasuji M, et al. C-C chemokine receptor 2 (CCR2) deficiency improves bleomycin-induced pulmonary fibrosis by attenuation of both macrophage infiltration and production of macrophage-derived matrix metalloproteinases. *J Pathol.* 2004;204:594-604. (IHC-FS)
8. Hasegawa T, Iwanaga K, Hultquist DE, Liao H, Visovatti SH, Pinsky DJ. Suppression of nitrosative and oxidative stress to reduce cardiac allograft vasculopathy. *Am J Physiol Heart Circ Physiol.* 2009;296:H1007-16. (IHC-FS)
9. Miyazaki Y, Kaikita K, Endo M, Horio E, Miura M, Tsujita K, et al. C/EBP homologous protein deficiency attenuates myocardial reperfusion injury by inhibiting myocardial apoptosis and inflammation. *Arterioscler Thromb Vasc Biol.* 2011;31:1124-32. (IHC-FS)
10. Hasegawa T, Okada K, Okita Y, Pinsky DJ. Antioxidant properties of pioglitazone limit nicotinamide adenine dinucleotide phosphate hydrogen oxidase and augment superoxide dismutase activity in cardiac allotransplantation. *J Heart Lung Transplant.* 2011;30:1186-96. (IHC-FS)
11. Scarff KL, Ung KS, Nandurkar H, Crack PJ, Bird CH, Bird PI. Targeted disruption of SPI3/Serpinb6 does not result in developmental or growth defects, leukocyte dysfunction, or susceptibility to stroke. *Mol Cell Biol.* 2004;24:4075-82. (FC)
12. De Cesare M, Calcaterra C, Pratesi G, Gatti L, Zunino F, Ménard S, et al. Eradication of ovarian tumor xenografts by locoregional administration of targeted immunotherapy. *Clin Cancer Res.* 2008;14:5512-8. (FC)

Spectral Red[®] is a registered trademark of Southern Biotechnology Associates, Inc.

Spectral Red[®] is a PE/CY5 tandem conjugate.

Cy[®] is a registered trademark of GE Healthcare.

Alexa Fluor[®] 488, 647, 700 and Pacific Blue[™] are provided under an agreement between Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corporation), and Southern Biotechnology Associates, Inc., and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications, and corresponding non-U.S. equivalents, owned by Molecular Probes, Inc. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Molecular Probes, Inc., Business Development, 29851 Willow Creek Road, Eugene, OR 97402, USA, Tel: (541) 465-8300. Fax: (541) 335-0504.