




Goat F(ab')₂ Anti-Human IgM

Cat. No.	Format	Size
2022-01	Purified (UNLB)	0.5 mg
2022-02	Fluorescein (FITC)	0.5 mg
2022-05	Horseradish Peroxidase (HRP)	1.0 mL
2022-08	Biotin (BIOT)	0.5 mg
2022-09	R-phycoerythrin (PE)	0.25 mg
2022-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg
2022-30	Alexa Fluor 488 [®] (AF488)	0.5 mg
2022-31	Alexa Fluor 647 [®] (AF647)	0.5 mg
2022-32	Alexa Fluor 555 [®] (AF555)	0.5 mg



Human peripheral blood lymphocytes were stained with Goat F(ab')₂ Anti-Human IgM-PE (SB Cat. 2022-09) and Mouse Anti-Human CD19-FITC (SB Cat. No. 9340-02).

Description

Specificity	Reacts with the heavy chain of human IgM
Source	Pepsin digest of Goat Anti-Human IgM (SB Cat. No. 2020)
Cross Adsorption	Human IgG and IgA; may react with IgM from other species

Applications

Quality tested applications include –

ELISA¹
FLISA
FC^{3,4}

Other referenced applications include –

ICC¹⁵
ELISPOT²
Stim⁵⁻¹⁴

Working Dilutions

ELISA	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC, AF488, and AF555 conjugates	1:100 – 1:400
	PE and AF647 conjugates	≤ 1 µg/mL
Flow Cytometry	FITC, BIOT, and AF488 conjugates	≤ 1 µg/10 ⁶ cells
	PE and AF647 conjugates	≤ 0.1 µg/10 ⁶ cells
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL	
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: info@southernbiotech.com • Website: www.southernbiotech.com

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC), biotin (BIOT), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 0.5 mg in 1.0 mL of PBS/Na₃. Store at 2-8°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.25 mg in 1.0 mL of PBS/Na₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of PBS. **Aliquot and store at or below -20°C.**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Suryani S, Fulcher DA, Santner-Nanan B, Nanan R, Wong M, Shaw PJ, et al. Differential expression of CD21 identifies developmentally and functionally distinct subsets of human transitional B cells. *Blood*. 2010;115:519-29. (ELISA)
2. Sira MM, Yoshida T, Takeuchi M, Kashiwayama Y, Futatani T, Kanegane H, et al. A novel immunoregulatory protein in human colostrum, syntenin-1, for promoting the development of IgA-producing cells from cord blood B cells. *Int Immunol*. 2009;21:1013-23. (ELISPOT)
3. White CA, Pone EJ, Lam T, Tat C, Hayama KL, Li G, et al. Histone deacetylase inhibitors upregulate B cell microRNAs that silence AID and Blimp-1 expression for epigenetic modulation of antibody and autoantibody responses. *J Immunol*. 2014;193:5933-50. (FC)
4. Westera L, van Hoeven V, Drylewicz J, Spierenburg G, van Velzen JF, de Boer RJ, et al. Lymphocyte maintenance during healthy aging requires no substantial alterations in cellular turnover. *Aging Cell*. 2015 Jan 28. doi: 10.1111/ace.12311. [Epub ahead of print]. (FC)
5. Petlickovski A, Laurenti L, Li X, Marietti S, Chiusolo P, Sica S, et al. Sustained signaling through the B-cell receptor induces Mcl-1 and promotes survival of chronic lymphocytic leukemia B cells. *Blood*. 2005;105:4820-7. (Stim)
6. Gobessi S, Laurenti L, Longo PG, Sica S, Leone G, Efremov DG. ZAP-70 enhances B-cell-receptor signaling despite absent or inefficient tyrosine kinase activation in chronic lymphocytic leukemia and lymphoma B cells. *Blood*. 2007;109:2032-9. (Stim)
7. Arechiga AF, Habib T, He Y, Zhang X, Zhang Z, Funk A, et al. Cutting edge: the PTPN22 allelic variant associated with autoimmunity impairs B cell signaling. *J Immunol*. 2009;182:3343-7. (Stim)
8. Chen K, Xu W, Wilson M, He B, Miller NW, Bengten E, et al. Immunoglobulin D enhances immune surveillance by activating antimicrobial, pro-inflammatory and B cell-stimulating programs in basophils. *Nat Immunol*. 2009;10:889-98. (Stim)
9. Suljagic M, Laurenti L, Tamani M, Alam M, Malek SN, Efremov DG. Reduced expression of the tumor suppressor PHLPP1 enhances the antiapoptotic B-cell receptor signal in chronic lymphocytic leukemia B-cells. *Leukemia*. 2010;24:2063-71. (Stim)
10. Cheng S, Coffey G, Zhang XH, Shaknovich R, Song Z, Lu P, et al. SYK inhibition and response prediction in diffuse large B-cell lymphoma. *Blood*. 2011;118:6342-52. (Stim)
11. Saiya-Cork K, Collins R, Parkin B, Ouillette P, Kuizon E, Kujawski L, et al. A pathobiological role of the insulin receptor in chronic lymphocytic leukemia. *Clin Cancer Res*. 2011;17:2679-92. (Stim)
12. Rankin AL, Seth N, Keegan S, Andreyeva T, Cook TA, Edmonds J, et al. Selective inhibition of BTK prevents murine lupus and antibody-mediated glomerulonephritis. *J Immunol*. 2013;191:4540-50. (Stim)
13. Jackson LJ, Selva S, Niedzielko T, Vollmer T. B cell receptor recognition of glatiramer acetate is required for efficacy through antigen presentation and cytokine production. *J Clin Cell Immunol*. 2014;5:185. (Stim)
14. Macmillan H, Strohmman MJ, Ayyangar S, Jiang W, Rajasekaran N, Spura A, et al. The MHC class II cofactor HLA-DM interacts with Ig in B cells. *J Immunol*. 2014;193:2641-50. (Stim)
15. Lloyd KA, Wang J, Urban BC, Czajkowsky DM, Pleass RJ. Glycan-independent binding and internalization of human IgM to FCμR, its cognate cellular receptor. *Sci Rep*. 2017;7:42989. (ICC)

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.