




Goat Anti-Guinea Pig IgG(H+L)

Cat. No.	Format	Size
6090-01	Purified (UNLB)	1.0 mg
6090-02	Fluorescein (FITC)	1.0 mg
6090-04	Alkaline Phosphatase (AP)	1.0 mL
6090-05	Horseradish Peroxidase (HRP)	1.0 mL
6090-08	Biotin (BIOT)	1.0 mg



ELISA plate was coated with purified guinea pig IgG. Immunoglobulin was detected with Goat Anti-Guinea Pig IgG(H+L)-BIOT (SB Cat. No. 6090-08) followed by Streptavidin-HRP (SB Cat. No. 7100-05).

Description

Specificity	Reacts with the heavy and light chains of guinea pig IgG
Source	Pooled antisera from goats hyperimmunized with guinea pig IgG
Cross Adsorption	None; may react with immunoglobulins from other species and the light chains of other guinea pig immunoglobulins
Purification	Affinity chromatography on guinea pig IgG covalently linked to agarose

Applications

Quality tested applications include –

ELISA¹⁻⁹

FLISA

Other referenced applications include –

IHC-FS^{10,11}

IHC-PS¹²⁻¹⁴

ICC^{15,16}

WB¹⁷⁻²¹

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC conjugate	1:200 – 1:400

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: info@southernbiotech.com • Website: www.southernbiotech.com

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Coëffier E, Excler J, Kieny MP, Meignier B, Moste C, Tartaglia J, et al. Restricted specificity of anti-V3 antibodies induced in humans by HIV candidate vaccines. *AIDS Res Hum Retroviruses*. 1997;13:1471-85. (ELISA)
2. McCracken AA, Werner ED, Powell MJ, Kruse KB, Brodsky JL. Differential fates of invertase mutants in the yeast endoplasmic reticulum. *Yeast*. 2000;16:49-55. (ELISA)
3. Pilorz V, Jäckel M, Knudsen K, Trillmich F. The cost of a specific immune response in young guinea pigs. *Physiol Behav*. 2005;85:205-11. (ELISA)
4. Ryder AB, Tollefson SJ, Podsiad AB, Johnson JE, Williams JV. Soluble recombinant human metapneumovirus G protein is immunogenic but not protective. *Vaccine*. 2010;28:4145-52. (ELISA, WB)
5. de Jonge MI, Keizer SA, el Moussaoui HM, van Dorsten L, Azzawi R, van Zuilekom HI, et al. A novel guinea pig model of Chlamydia trachomatis genital tract infection. *Vaccine*. 2011;29:5594-6001. (ELISA)
6. Velasquez LS, Shira S, Berta AN, Kilbourne J, Medi BM, Tizard I, et al. Intranasal delivery of Norwalk virus-like particles formulated in an in situ gelling, dry powder vaccine. *Vaccine*. 2011;29:5221-31. (ELISA)
7. Schiött Å, Larsson K, Manniche S, Kalliomaäki S, Heydenreich AV, Dalsgaard K, et al. Posintro™-HBsAg, a modified ISCOM including HBsAg, induces strong cellular and humoral responses. *Int J Pharm*. 2011;414:312-20. (ELISA)
8. Baldwin SL, Bertholet S, Reese VA, Ching LK, Reed SG, Coler RN. The importance of adjuvant formulation in the development of a tuberculosis vaccine. *J Immunol*. 2012;188:2189-97. (ELISA)
9. Rebollar PG, Dal Bosco A, Millán P, Cardinali R, Brecchia G, Sylla L, et al. Ovulating induction methods in rabbit does: the pituitary and ovarian responses. *Theriogenology*. 2012;77:292-8. (ELISA)
10. Yamada K, Fukaya M, Shibata T, Kurihara H, Tanaka K, Inoue Y, et al. Dynamic transformation of Bergmann glial fibers proceeds in correlation with dendritic outgrowth and synapse formation of cerebellar Purkinje cells. *J Comp Neurol*. 2000;418:106-20. (IHC-FS)
11. Kraus LM, Gaber L, Handorf CR, Marti H, Kraus AP. Carbamylation of glomerular and tubular proteins in patients with kidney failure: a potential mechanism of ongoing renal damage. *Swiss Med Wkly*. 2001;131:139-45. (IHC-FS)
12. Fukaya M, Yamada K, Nagashima M, Tanaka K, Watanabe M. Down-regulated expression of glutamate transporter GLAST in Purkinje cell-associated astrocytes of reeler and weaver mutant cerebella. *Neurosci Res*. 1999;34:165-75. (IHC-PS)
13. Klymiuk N, van Buerck L, Bähr A, Offers M, Kessler B, Wuensch A, et al. Xenografted islet clusters from INSLEA29Y transgenic pigs rescue diabetes and prevent immune rejection in humanized mice. *Diabetes*. 2012;61:1527-32. (IHC-PS)
14. Streckel E, Braun-Reichhart C, Herbach N, Dahlhoff M, Kessler B, Blutke A. Effects of the glucagon-like peptide-1 receptor agonist liraglutide in juvenile transgenic pigs modeling a pre-diabetic condition. *J Transl Med*. 2015;13:73. (IHC-PS)
15. Yao Q, Compans RW. Filamentous particle formation by human parainfluenza virus type 2. *J Gen Virol*. 2000;81:1305-12. (ICC)
16. Cseke G, Maginnis MS, Cox RG, Tollefson SJ, Podsiad AB, Wright DW, et al. Integrin αvβ1 promotes infection by human metapneumovirus. *Proc Natl Acad Sci USA*. 2009;106:1566-71. (ICC)
17. Vasilakos JP, Michael JG. Herpes simplex virus class I-restricted peptide induces cytotoxic T lymphocytes in vivo independent of CD4⁺ T cells. *J Immunol*. 1993;150:2346-55. (WB)
18. King MC, Raposo G, Lemmon MA. Inhibition of nuclear import and cell-cycle progression by mutated forms of the dynamin-like GTPase MxB. *Proc Natl Acad Sci USA*. 2004;101:8957-62. (WB)
19. Dalen KT, Ulven SM, Arntsen BM, Solaas K, Nebb HI. PPARα activators and fasting induce the expression of adipose differentiation-related protein in liver. *J Lipid Res*. 2006;47:931-43. (WB)
20. Cseke G, Wright DW, Tollefson SJ, Johnson JE, Crowe JE Jr, Williams JV. Human metapneumovirus fusion protein vaccines that are immunogenic and protective in cotton rats. *J Virol*. 2007;81:698-707. (WB)
21. Nobre RJ, Herráez-Hernández E, Fei J, Langbein L, Kaden S, Gröne H, et al. E7 oncoprotein of novel human papillomavirus type 108 lacking the E6 gene induces dysplasia in organotypic keratinocyte cultures. *J Virol*. 2009;83:2907-16. (WB)