




Mouse Anti-Chicken Lambda

| Cat. No. | Format | Size |
|----------|----------------------|--------|
| 8340-01 | Purified (UNLB) | 0.5 mg |
| 8340-02 | Fluorescein (FITC) | 0.5 mg |
| 8340-08 | Biotin (BIOT) | 0.5 mg |
| 8340-09 | R-phycoerythrin (PE) | 0.1 mg |



Chicken peripheral blood lymphocytes were stained with Mouse Anti-Chicken Lambda-PE (SB Cat. No. 8340-09) and Mouse Anti-Chicken CD3-FITC (SB Cat. No. 8200-02).

Overview

| | |
|--------------------------|--|
| Clone | L-1 |
| Isotype | Mouse (BALB/c) IgG ₁ κ |
| Immunogen | Affinity purified chicken Ig or isolated lymphocytes |
| Specificity | Chicken Lambda; Mr 28 kDa |
| Alternate Name(s) | N/A |

Applications

FC – Quality tested ³⁻⁸
 IHC-FS – Reported in literature ²
 ICC – Reported in literature ¹
 IP – Reported in literature ¹
 WB – Reported in literature ³

Working Dilutions

| | | |
|---------------------------|--|------------------------------|
| Flow Cytometry | FITC, BIOT, and PE conjugates For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL | ≤ 1 µg/10 ⁶ cells |
| Other Applications | Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need. | |

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Chen CH, Lehmeyer JE, Cooper MD. Evidence for an IgD homologue on chicken lymphocytes. J Immunol. 1982;129:2580-5. (Immunogen, IP, ICC)
2. O'Halloran EK, Oesterle EC. Characterization of leukocyte subtypes in chicken inner ear sensory epithelia. J Comp Neurol. 2004;475:340-60. (IHC-FS)
3. Luo J, Zhang H, Teng M, Fan J, You L, Xiao Z, et al. Surface IgM on DT40 cells may be a component of the putative receptor complex responsible for the binding of infectious bursal disease virus. Avian Pathol. 2010;39:359-65. (WB, FC)
4. Wang YW, Cherian G, Sunwoo HH, Sim JS. Dietary polyunsaturated fatty acids significantly affect laying hen lymphocyte proliferation and immunoglobulin G concentration in serum and egg yolk. Can J Anim Sci. 2000;80:597-604. (FC)
5. Islam AF, Wong CW, Walkden-Brown SW, Colditz IG, Arzey KE, Groves PJ. Immunosuppressive effects of Marek's disease virus (MDV) and herpesvirus of turkeys (HVT) in broiler chickens and the protective effect of HVT vaccination against MDV challenge. Avian Pathol. 2002;31:449-61. (FC)
6. Nera K, Alinikula J, Terho P, Narvi E, Törnquist K, Kurosaki T, et al. Ikaros has a crucial role in regulation of B cell receptor signaling. Eur J Immunol. 2006;36:516-25. (FC)
7. Viertlboeck BC, Schweinsberg S, Hanczaruk MA, Schmitt R, Du Pasquier L, Herberg FW, et al. The chicken leukocyte receptor complex encodes a primordial, activating, high-affinity IgY Fc receptor. Proc Natl Acad Sci USA. 2007;104:11718-23. (FC)
8. Schusser B, Collarini EJ, Yi H, Izquierdo SM, Fesler J, Pedersen D, et al. Immunoglobulin knockout chickens via efficient homologous recombination in primordial germ cells. Proc Natl Acad Sci USA. 2013;110:20170-5. (FC)