




## Goat Anti-Mouse IgE

Cat. No.	Format	Size
1110-01	Purified (UNLB)	1.0 mg
1110-02	Fluorescein (FITC)	1.0 mg
1110-04	Alkaline Phosphatase (AP)	1.0 mL
1110-05	Horseradish Peroxidase (HRP)	1.0 mL
1110-08	Biotin (BIOT)	1.0 mg
1110-09	R-phycoerythrin (PE)	0.5 mg



FLISA plate was coated with purified mouse IgE, IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>, IgG<sub>3</sub>, IgM, and IgA. Immunoglobulins were detected with serially diluted Goat Anti-Mouse IgE-FITC (SB Cat. No. 1110-02).

### Description

<b>Specificity</b>	Reacts with the heavy chain of mouse IgE
<b>Source</b>	Pooled antisera from goats hyperimmunized with mouse IgE
<b>Cross Adsorption</b>	Mouse IgG <sub>1</sub> , IgG <sub>2a</sub> , IgG <sub>2b</sub> , IgG <sub>3</sub> , IgM, and IgA; may react with immunoglobulins from other species
<b>Purification</b>	Affinity chromatography on mouse IgE covalently linked to agarose

### Applications

Quality tested applications include –

ELISA<sup>1-17</sup>  
FLISA

Other referenced applications include –

FC<sup>18</sup>  
IHC<sup>19</sup>  
WB<sup>20,21</sup>  
IP<sup>22</sup>

### Working Dilutions

<b>ELISA</b>	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
<b>FLISA</b>	FITC conjugate	1:100 – 1:400
	PE conjugate	≤ 1 µg/mL

**Other Applications** Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: [info@southernbiotech.com](mailto:info@southernbiotech.com) • Website: [www.southernbiotech.com](http://www.southernbiotech.com)

## Handling and Storage

---

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 1.0 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl<sub>2</sub>/50% glycerol, pH 8.0, containing NaN<sub>3</sub> as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

---

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

## References

---

1. Costa-Pinto FA, Basso AS, Britto LR, Malucelli BE, Russo M. Avoidance behavior and neural correlates of allergen exposure in a murine model of asthma. *Brain Behav Immun.* 2005;19:52-60. (ELISA)
2. Andrade MC, Menezes JS, Cassali GD, Martins-Filho OA, Cara DC, Faria AM. Alcohol-induced gastritis prevents oral tolerance induction in mice. *Clin Exp Immunol.* 2006;146:312-22. (ELISA)
3. Prescott VE, Forbes E, Foster PS, Matthaei K, Hogan SP. Mechanistic analysis of experimental food allergen-induced cutaneous reactions. *J Leukoc Biol.* 2006;80:258-66. (ELISA)
4. Annaboldi PM, Roth-Walter F, Mayer L. Suppression of Th1 and Th17, but not Th2, responses in a CD8<sup>+</sup> T cell mediated model of oral tolerance. *Mucosal Immunol.* 2009;2:427-38. (ELISA)
5. Pavoglio S, Puddington L, Rafti E, Matson AP. FcRn-mediated intestinal absorption of IgG anti-IgE/IgE immune complexes in mice. *Clin Exp Immunol.* 2012;42:1791-800. (ELISA)
6. Kawabe Y, Hayashida Y, Numata K, Harada S, Hayashida Y, Ito A, et al. Oral immunotherapy for pollen allergy using T-cell epitope-containing egg white derived from genetically manipulated chickens. *PLoS One.* 2012;7(10):e48512. (ELISA)
7. Kin NW, Stefanov EK, Dizon BL, Kearney JF. Antibodies generated against conserved antigens expressed by bacteria and allergen-bearing fungi suppress airway disease. *J Immunol.* 2012;189:2246-56. (ELISA)
8. Lochmatter C, Schneider CL, Ingram K, Keiser J, Schifferli JA. Schistosoma mansoni tetraspanning orphan receptor (SmTOR): a new vaccine candidate against schistosomiasis. *Clin Exp Immunol.* 2012;170:342-57. (ELISA)
9. Durham CG, Schwiebert LM, Lorenz RG. Use of the cockroach antigen model of acute asthma to determine the immunomodulatory role of early exposure to gastrointestinal infection. *Methods Mol Biol.* 2013;1032:271-86. (ELISA)
10. Vukcevic M, Zorzato F, Keck S, Tsakiris DA, Keiser J, Maizels RM, et al. Gain of function in the immune system caused by a ryanodine receptor 1 mutation. *J Cell Sci.* 2013;126:3485-92. (ELISA)
11. Brachs S, Turqueti-Neves A, Stein M, Reimer D, Brachvogel B, Bösl M, et al. Swiprosin-1/EFhd2 limits germinal center responses and humoral type 2 immunity. *Eur J Immunol.* 2014;44:3206-19. (ELISA)
12. Zhou W, Goleniewska K, Zhang J, Dulek DE, Toki S, Lotz MT, et al. Cyclooxygenase inhibition abrogates aeroallergen-induced immune tolerance by suppressing prostaglandin I<sub>2</sub> receptor signaling. *J Allergy Clin Immunol.* 2014;134:698-705. (ELISA)
13. Joshi VB, Adamcakova-Dodd A, Jing X, Wongrakpanich A, Gibson-Corley KN, Thorne PS, et al. Development of a poly (lactic-co-glycolic acid) particle vaccine to protect against house dust mite induced allergy. *AAPS J.* 2014;16:975-85. (ELISA)
14. Sun R, Tang X, Yao H, Hong S, Yang Y, Kou W, et al. Establishment of a new animal model of allergic rhinitis with biphasic sneezing by intranasal sensitization with Staphylococcal enterotoxin B. *Exp Ther Med.* 2015;10:407-12. (ELISA)
15. Bracken SJ, Adams AJ, Szczepanek SM, Ehsan M, Natarajan P, Guernsey LA, et al. Long-term exposure to house dust mite leads to the suppression of allergic airway disease despite persistent lung inflammation. *Int Arch Allergy Immunol.* 2015;166:243-58. (ELISA)
16. Patel PS, Kearney JF. Neonatal exposure to pneumococcal phosphorylcholine modulates the development of house dust mite allergy during adult life. *J Immunol.* 2015;194:5838-50. (ELISA)
17. Chen X, Lin J, Pan T, Cao M, Shi C, Cai Q, et al. Investigation of the hazardous substance causing crayfish-induced rhabdomyolysis via a mouse model, a hemolysis assay, and a cytotoxicity assay. *Fish Sci.* 2015;81:551-8. (ELISA)
18. Vogel L, Lüttkopf D, Hatahet L, Hausteiner D, Vieths S. Development of a functional in vitro assay as a novel tool for the standardization of allergen extracts in the human system. *Allergy.* 2005;60:1021-8. (FC)
19. Samarasinghe AE, Hoselton SA, Schuh JM. The absence of VPAC<sub>2</sub> leads to aberrant antibody production in Aspergillus fumigatus sensitized and challenged mice. *Peptides.* 2011;32:131-7. (IHC)
20. Burns TA, Westerman T, Nuovo GJ, Watts MR, Pettigrew A, Yin C, et al. Role of oxidative tissue injury in the pathophysiology of experimentally induced equine laminitis: a comparison of 2 models. *J Vet Intern Med.* 2011;25:540-8. (WB)
21. Nieuwenhuizen NE, Meter JM, Horsnell WG, Hoving JC, Fick L, Sharp MF, et al. A cross-reactive monoclonal antibody to nematode haemoglobin enhances protective immune responses to Nippostrongylus brasiliensis. *PLoS Negl Trop Dis.* 2013;7(8):e2395. (WB)
22. Hazenbos WL, Wu P, Eastham-Anderson J, Kinoshita T, Brown EJ. Impaired FcεRI stability, signaling, and effector functions in murine mast cells lacking glycosylphosphatidylinositol-anchored proteins. *Blood.* 2011;118:4377-83. (IP)