




## Mouse Anti-Mouse Ly-49I

Cat. No.	Format	Size
1802-01	Purified (UNLB)	0.5 mg
1802-02	Fluorescein (FITC)	0.5 mg
1802-08	Biotin (BIOT)	0.5 mg
1802-09	R-phycoerythrin (PE)	0.1 mg



C57BL/6 mouse splenocytes were stained with Mouse Anti-Mouse Ly-49I-BIOT (SB Cat. 1802-08) and Mouse Anti-Mouse NK1.1-PE (SB Cat. No. 1805-09) followed by Streptavidin-FITC (SB Cat. No. 7100-02).

### Overview

<b>Clone</b>	YLI-90
<b>Isotype</b>	Mouse (BALB/c) IgG <sub>1κ</sub>
<b>Immunogen</b>	CHO-K1 cells transfected with B6 allele of Ly49I gene
<b>Specificity</b>	Mouse Ly-49I
<b>Alternate Name(s)</b>	Ly-49

### Description

The monoclonal antibody YLI-90 reacts with the B6 allele of Ly49I, a member of the lectin-like homodimeric Ly49 family of cell surface receptors. Mouse Ly49 receptors exhibit allelic specificity for MHC class I Ia molecules and are thought to serve to prevent natural killer (NK) cells from attacking normal cells while allowing them to attack infected or transformed cells in which class I molecules have been down-regulated. These inhibitory receptors are also expressed on subpopulations of mouse CD8<sup>+</sup> T cells. The YLI-90 antibody stains ~45% of NK1.1<sup>+</sup>CD3<sup>-</sup> splenocytes from C57BL/6 mice.

### Applications

FC – Quality tested <sup>1,2</sup>

### Working Dilutions

<b>Flow Cytometry</b>	Purified (UNLB) antibody	≤ 1 μg/10 <sup>6</sup> cells
	FITC and BIOT conjugates	≤ 1 μg/10 <sup>6</sup> cells
	PE conjugate	≤ 0.2 μg/10 <sup>6</sup> cells

For flow cytometry, the suggested use of these reagents is in a final volume of 100 μL

**Other Applications** Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: [info@southernbiotech.com](mailto:info@southernbiotech.com) • Website: [www.southernbiotech.com](http://www.southernbiotech.com)

## Handling and Storage

---

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

---

Some reagents contain sodium azide. Please refer to product specific SDS.

## References

---

1. Coles MC, McMahon CW, Takizawa H, Raulet DH. Memory CD8 T lymphocytes express inhibitory MHC-specific Ly49 receptors. *Eur J Immunol.* 2000;30:236-44. (Immunogen, FC)
2. Corral L, Takizawa H, Hanke T, Jamieson AM, Raulet DH. A new monoclonal antibody reactive with several Ly49 NK cell receptors mediates redirected lysis of target cells. *Hybridoma.* 1999;18:359-66. (FC)

Important Licensing Information - These products may be covered by one or more Limited Use Label Licenses. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses.