




Mouse Anti-Human IgE Fc

Cat. No.	Format	Size
9160-01	Purified (UNLB)	0.5 mg
9160-02	Fluorescein (FITC)	0.5 mg
9160-04	Alkaline Phosphatase (AP)	1.0 mL
9160-05	Horseradish Peroxidase (HRP)	1.0 mL
9160-08	Biotin (BIOT)	0.5 mg
9160-09	R-phycoerythrin (PE)	0.1 mg
9160-31	Alexa Fluor® 647 (AF647)	0.1 mg
9160-32	Alexa Fluor® 555 (AF555)	0.1 mg



ELISA plate was coated with purified human IgG, IgM, IgA, and IgE. Immunoglobulins were detected with serially diluted Mouse Anti-Human IgE Fc-HRP (SB Cat. No. 9160-05).

Overview

Clone	B3102E8
Isotype	Mouse IgG _{1κ}
Immunogen	Unknown
Specificity	Human IgE Fc; Mr 188 kDa

Applications

ELISA – Quality tested ¹⁻³
 FLISA – Quality tested
 FC – Reported in literature ¹⁶
 IHC-FS – Reported in literature ⁴⁻¹³
 ICC – Reported in literature ¹²
 WB – Reported in literature ^{2,3,12,14,15}

Working Dilutions

ELISA	AP conjugate	1:500 – 1:1,000
	HRP conjugate	1:1,000 – 1:2,000
	BIOT conjugate	1:5,000 – 1:10,000
FLISA	FITC and AF555 conjugates	1:100 – 1:400
	PE and AF647 conjugates	≤ 1 µg/mL

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL in a stock solution of 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL in a stock solution of 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Alexa Fluor® 555 (AF555) and Alexa Fluor® 647 (AF647) conjugates are supplied as 0.1 mg in 0.2 mL of PBS/NaN₃. Store at 2-8°C.
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Xue L, Johnson R, Gorovits B. Prevalence and isotypic complexity of the anti-Chinese hamster ovary host cell protein antibodies in normal human serum. *AAPS J.* 2010;12:98-106. (ELISA)
2. Goodman RE, Panda R, Ariyaratna H. Evaluation of endogenous allergens for the safety evaluation of genetically engineered food crops: review of potential risks, test methods, examples and relevance. *J Agric Food Chem.* 2013;61:8317-32. (ELISA, WB)
3. Berkner H, Seutter von Loetzen C, Hartl M, Randow S, Gubesch M, Vogel L, et al. Enlarging the toolbox for allergen epitope definition with an allergen-type model protein. *PLoS One.* 2014;9(10):e111691. (ELISA, WB)
4. Meding B, Ahonen K, Brisman J, Hedblad M, Ståhl A, Järholm B, et al. Late skin-prick-test reactions to malted wheat. Clinical observations and immunohistochemical characterization. *Allergy.* 1998;53:282-8. (IHC-FS)
5. Tengvall Linder M, Johansson C, Scheynius A, Wahlgren C. Positive atopy patch test reactions to *Pityrosporum orbiculare* in atopic dermatitis patients. *Clin Exp Allergy.* 2000;30:122-31. (IHC-FS)
6. Åström K, Magnusson CG, Papatziamos G, Hemlin C, Scheynius A, van der Ploeg I, et al. Analysis of epsilon germline transcripts and IL-4 mRNA expression in the adenoids suggests local IgE switching. *Allergy.* 2000;55:1049-55. (IHC-FS)
7. Sverremark Ekström E, Nilsson C, Holmlund U, van der Ploeg I, Sandstedt B, Lilja G, et al. IgE is expressed on, but not produced by, fetal cells in the human placenta irrespective of maternal atopy. *Clin Exp Immunol.* 2002;127:274-82. (IHC-FS)
8. Nivenius E, Montan PG, Chryssanthou E, Jung K, van Hage-Hamsten M, van der Ploeg I. No apparent association between periorbital and ocular microcolonization and the degree of inflammation in patients with atopic keratoconjunctivitis. *Clin Exp Allergy.* 2004;34:725-30. (IHC-FS)
9. Rindsjö E, Varli IH, Ofori MF, Lundquist M, Holmlund U, Papadogiannakis N, et al. Presence of IgE⁺ cells in human placenta is independent of malaria infection or chorioamnionitis. *Clin Exp Immunol.* 2006;144:204-11. (IHC-FS)
10. Forshammar J, Isaksson S, Strid H, Stotzer P, Sjövall H, Simrén M, et al. A pilot study of colonic B cell pattern in irritable bowel syndrome. *Scand J Gastroenterol.* 2008;43:1461-66. (IHC-FS)
11. Spazierer D, Skvara H, Dawid M, Fallahi N, Gruber K, Rose K, et al. T helper 2 biased de novo immune response to Keyhole Limpet Hemocyanin in humans. *Clin Exp Allergy.* 2009;39:999-1008. (IHC-FS)
12. Altrichter S, Kriehuber E, Moser J, Valenta R, Kopp T, Stingl G. Serum IgE autoantibodies target keratinocytes in patients with atopic dermatitis. *J Invest Dermatol.* 2008;128:2232-9. (IHC-FS, ICC, WB)
13. Joerink M, Rindsjö E, Stenius F, Alm J, Lilja G, Grönlund H, et al. Evidence for allergen-specific IgE of maternal origin in human placenta. *Allergy.* 2009;64:905-12. (IHC-FS)
14. Gómez-Ollés S, Cruz MJ, Renström A, Doekes G, Morell F, Rodrigo MJ. An amplified sandwich EIA for the measurement of soy aeroallergens. *Clin Exp Allergy.* 2006;36:1176-83. (WB)
15. Venkatachalam M, Teuber SS, Peterson WR, Roux KH, Sathe SK. Antigenic stability of pecan [*Carya illinoensis* (Wangenh.) K. Koch] proteins: effects of thermal treatments and in vitro digestion. *J Agric Food Chem.* 2006;54:1449-58. (WB)
16. Buentke E, Zargari A, Heffler LC, Avila-Cariño J, Savolainen J, Scheynius A. Uptake of the yeast *Malassezia furfur* and its allergenic components by human immature CD1a⁺ dendritic cells. *Clin Exp Allergy.* 2000;30:1759-70. (FC)

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

TB9160
05-Dec-21

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • **Fax:** 205.945.8768

Email: info@southernbiotech.com • **Website:** www.southernbiotech.com